

Clark  
1920

## THE MOLLUSKS

### THE UNIONIDÆ

During the study of Lake Maxinkuckee considerable attention was given to the mollusks, particularly the freshwater mussels or Unionidæ. This was justified by the rapid and astonishing development of the pearl button industry in America which is dependent upon the shells of mussels for its raw material. The recent discovery by Lefevre and Curtis of methods whereby commercially valuable species of mussels are now successfully propagated artificially, adds special interest to these mollusks. Recent studies and discoveries relating to the formation and artificial production of pearls in freshwater mussels may also be mentioned as showing the importance of careful study of the taxonomy and life history of the Unionidæ.

### LAKES AND PONDS AS THE HOME OF MUSSELS

Generally speaking, lakes and ponds are not so well suited to the growth and development of mussels as rivers are; the species of lake or pond mussels are comparatively few, and the individuals usually somewhat dwarfed. Of about 84 species of mussels reported for the State of Indiana, only about 24 are found in lakes, not all of these in any one lake, and several of them but rarely in any. Of the 24 species occasionally found in lakes, but five are reported only in lakes, and only three or four of the species common to both lakes and rivers seem to prefer lakes.

In rivers, the essential feature favorable to the development of mussels is the current; and in the rivers the mussel beds reach their best development in the riffles, where the current is strongest. The importance of the current to the well-being of the mussels is indicated by the position these mollusks naturally assume in the beds, the inhalent and exhalent apertures of the creatures being directed up-stream against the current. The importance of the current is not merely as a bringer of food; examinations show that the mussels of the plankton-rich lakes and ponds usually contain more food material than those of rivers. The current gives the river-mussels the advantage of a constant change of water, which means a more abundant supply of oxygen, and doubtless a more varied supply of mineral matter, from the various sorts of soil through which the river flows.

The current is also probably of considerable importance in assisting in the fertilization of mussels, one of its functions being

the conveyance of sperm from mussels in upper portions of the bed to other mussels below. In places where there is no current, fertilization must be more largely a matter of chance.

Although the majority of species of mussels prefer a river where there is a good current, some are more fitted to the quieter parts of streams, or to ponds. These are chiefly thin-shelled species with weakly developed or undeveloped hinge-teeth, best represented by the genus *Anodonta*. In some places *Anodonta*s are known as pond-mussels, as distinguished from the heavier sorts or river-mussels.

The distinction between lakes and rivers is not constant in degree; we have all sorts of gradations from the extreme form of lake—isolated bodies without outlet—through lakes with relatively large, important outlets, to such lakes as are simply expansions of a river-bed, examples of the latter type being Lake Pepin, Minn., of the upper Mississippi, and the former English Lake in Indiana, an expansion of the Kankakee. As a usual thing, the more fluviatile a lake is, or the larger and more river-like its outlet, the more river-like will be its mussel fauna, both in abundance and species. In such lakes the mussels retain a vital continuity with the mussel beds of the river. In the less fluviatile lakes the mussels are more isolated, and there is more inbreeding. The large number (24) of lake-dwelling species recorded for Indiana is due to the fact that some of the lakes of Indiana are more or less fluviatile, and contain several species of river shells.

#### ORIGIN AND CHARACTER OF THE MAXINKUCKEE MUSSELS

Lake Maxinkuckee, having a long, narrow, winding and relatively unimportant outlet, is a representative of one of the less fluviatile types of lakes, forming a pretty well marked contrast to the various lakes cited above, and bearing a pretty close resemblance to the neighboring lakes, such as Twin Lakes, Pretty Lake, Bass Lake, etc.

The Maxinkuckee mussels are doubtless derived from ancestors brought up the Outlet from the Tippecanoe River by ascending fishes. It is doubtful whether any have been introduced by the numerous plants of fish in the lake, though such a thing is possible. During the various times the lake was visited, a few Tippecanoe River mussels were planted in the thoroughfare between the lakes, and a few Yellow River and Kankakee mussels were planted in the main lake.

The Outlet of Lake Maxinkuckee is now a narrow, shallow winding stream, straightened in places by ditching, and bordered

of the  
urrent,  
  
river  
quieter  
l spec-  
repre-  
as are  
orts or  
  
in de-  
orm of  
atively  
ions of  
Minn.,  
ndiana,  
fluvia-  
e more  
species.  
mussel  
re more  
r (24)  
act that  
contain

on each side by a flat sedgy plain which indicates the former breadth and importance of the stream. The colonization of the lake with mussels was probably effected chiefly during the period when the Outlet was a broad and relatively important stream. The situation has been carefully considered and seems to show that the mussels of the river and lake are isolated from each other and that there is no longer any vital connection between them. The strongest indication of the independence of the lake and river mussel faunas is the appearance of the Maxinkuckee mussels themselves; these are lake-mussels, easily distinguished for the most part from river mussels of the same species, and many of them are ditinguishable also from the mussels of the neighboring lakes.

ELS  
nd rela-  
the less  
trast to  
resem-  
y Lake,  
  
om an-  
r by as-  
roduced  
thing is  
l, a few  
fare be-  
mussels

The Tippecanoe River is fairly well supplied with mussels. Although the number of species is considerably fewer, and the size of the individuals is generally smaller, than that of the Wabash into which it flows, it compares very favorably with rivers of its size. At Delong, Ind., a short distance above the mouth of the Outlet of Lake Maxinkuckee, were obtained in one bed specimens representing 24 species of mussels or about twice the number of kinds found in Lake Maxinkuckee.

om an-  
r by as-  
roduced  
thing is  
l, a few  
fare be-  
mussels

Our knowledge of the extent and importance of migrations of fishes from the Tippecanoe River up to the lake and from the lake down to the river—a question which has a marked bearing upon the relationship of the mussel faunas—is not as complete as it should be, but indications are that they are not important or extensive. Inasmuch as the geographic distribution of a given species of mussel is coextensive with that of the species of fish which serves as its host, this question is worthy of careful consideration. There are several species of fishes of the Tippecanoe River (*Etheostoma camurum*, *Hadropterus evides*, *Hybopsis amblops*, etc.) which were not found either in the Outlet or in the lakes, and other species (*Hadropterus aspro*, *Ericymba buccata*, *Diplesion blennioides*) which have pushed halfway up the Outlet, but were found no further up.

om an-  
r by as-  
roduced  
thing is  
l, a few  
fare be-  
mussels

In this connection, the mussel fauna of the Outlet is worthy of consideration, and on various occasions, but especially on a trip down the Outlet September 30, 1907, particular attention was paid to this feature.

shallow  
bordered

The Outlet is not particularly well suited to the life and growth of mussels; the bottom is either a firm peaty soil or fine shifting sand; moreover, the course has been artificially changed in some places and the stream has naturally shortened its length in others by making cutoffs. In addition to this the mussel fauna of such

a narrow shallow stream would be the prey of muskrats, minks, etc., the entire length and width of the beds.

On the trip mentioned above, about a mile below Lost Lake a fine example of *Lampsilis iris* was found. This is the farthest up stream any species of mussel was obtained, and as this species is fairly common in both lakes and abundant in the Tippecanoe River, we have here the nearest approach to a continuous fauna. Some dead shells but no living examples of *Quadrula undulata* were found a little farther down. Farther down stream, from a quarter to half a mile, a short distance above the second cross-road south of the lake, was found a small mussel-bed of about 40 or 50 mussels, the great majority of which were *Quadrula undulata*. A few living *Lampsilis iris*, two dead *Symphynota compressa*, one living *Symphynota costata* (gravid), and a few dead shells of *Quadrula coccinea*, complete the list. Below this point no mussels were found until near where the Outlet joins the Tippecanoe. Here, a few rods up the Outlet, a fair bed of *Quadrula coccinea* was found. Of the five species of mussel found in the Outlet, only two, *L. iris* and *Q. coccinea*, are found in the lake, the latter but rarely. The form and general appearance of the *Q. undulata* of the Outlet is quite peculiar and they can be picked out at once in collections from the various rivers of the country. They are unusually elongate, in this respect representing some of the Tippecanoe mussels but differing from them in being thinner, and in having the furrows between the plicæ unusually deep and sharp. The costæ on the postero-dorsal slope are very marked, and the epidermis jet black. The umbones are considerably eroded.

#### DISTRIBUTION OF MUSSELS IN THE LAKE

In rivers, where there is a great variety of conditions, such as differences of current, bottom, etc., one finds the different species of mussels inhabiting different localities and different situations. In the lakes, where we have comparatively few species of mussels and not such important differences of environment, the distribution of the various species is much the same. The same conditions, such as rather shallow water and moderately firm bottom, are equally suitable for all. A few important exceptions may be noted; as for example, the less common species of the lake are often more or less local in distribution. The only well-marked bed of *Quadrula rubiginosa* in the lakes is in the Lost Lake mussel-bed below the Bardsley cottage, and this is the only place where *Lampsilis subrostrata* can be collected in any considerable numbers. *Lampsilis glans* has a marked preference for the shallow water at

the  
am  
and  
Poi  
tha  
was

2 to  
ma  
ies  
gra  
spe  
sho  
Aul  
low  
fair  
the  
nor  
The  
low  
litt  
Lal  
Bar  
thi

tio  
lak  
qui  
one  
live  
sio  
sho  
in  
in  
ser  
of  
fur  
mu  
sho  
the  
anc  
anc  
abc

the edge of the thoroughfare between the lakes; occasional examples can, however, be picked up almost anywhere along shore, and it appears to be increasing considerably along shore at Long Point. *Anodonta grandis footiana*, which can live in softer bottom than the other mussels, has a considerably wider distribution, and was dredged up in deeper water than any of the other mussels.

The mussels are to be found almost anywhere in water from 2 to 5 or 6 feet deep where the bottom is more or less sandy or marly. The beds are composed chiefly of the three principal species of the lake, *Lampsilis luteola*, *Unio gibbosus* and *Anodonta grandis footiana*, with the less common species sparsely interspersed. Especially good mussel beds occur at Long Point, along shore by Farrar's and McDonald's, by the Depot grounds, in Aubeenaubee Bay out from the Military Academy, and in the shallow water just beyond the mouth of Norris Inlet. Mussels are fairly well scattered from Long Point more or less continuously all the way southward to beyond Overmyer's hill, and from a little north of the ice-houses all the way around to the Military Academy. They are quite abundant in the neighborhood of Winfield's in shallow water, and occur scattered along the east side of the lake a little way out from shore. A good mussel bed is found in Lost Lake along the east shore, extending from a little south of the Bardsley cottage to where the bulrushes and water-lilies grow thickly in the soft black muck near shore.

*Movements*:—Closely connected with the question of distribution is that of movement. The greater number of mussels of the lake, especially in the deeper water, spend their lives in a state of quiescence. Young mussels appear to be more active than older ones. The mussels retain the power of locomotion during all their lives, but after they have got well settled down, they only occasionally use this power. The mussels of the shallow water near shore move out into deeper water at the approach of cold weather, in late autumn or early winter, and bury themselves more deeply in the sand. This movement is rather irregular and was not observed every year. It was strikingly manifest in the late autumn of 1913, when at one of the piers off Long Point a large number of furrows was observed heading straight into deep water with a mussel at the outer end of each. The return of the mussels to shore during the spring and summer was not observed. Many of them are probably washed shoreward by strong waves of the spring and summer storms, and some are carried shoreward by muskrats and dropped there. Occasional mussels were observed moving about in midwinter—even in rather deep water. During the

winter of 1900-1901 an example of *Lampsilis luteola*, in rather deep water in the vicinity of Winfield's, was observed to have moved about 18 inches within a few days. Its track could be distinctly seen through the clear ice.

As a result of the quiescence of the lake mussels, the posterior half or  $\frac{1}{3}$  of the shell, which projects up from the lake bottom, is usually covered by a thick marly concretion which appears to be a mixture of minute algæ and lime. This marly concretion grows concentrically, forming rounded nodules, its development increasing with the age and size of the shell. This concretion, though most abundant on shells, is not confined entirely to them, but grows also on rocks that have lain undisturbed on the bottom. When growing on shells, it adheres to them very closely; and upon being pried loose sometimes separates from them much as the matrix separates from a fossil, and leaves the epidermis of the mussel clean. In other cases it adheres more firmly and is difficult to scrape off clean. On this marly growth, colonies of Ophrydium, much the size, color, and general appearance of grapes with the skins removed, are often found growing, and in the cavities and interstices of the marl, a handsome little water-beetle, *Stenelmis undulatus* Blatchley, and its peculiar elongate black larvæ, live in considerable numbers but apparently have nothing to do with the mussels. Various species of hydrachnids, one of them strikingly handsome with its green body sprinkled with bright red dots, also live in the cavities of the marl, and offer some suggestion as to how the parasitic mite *Atax* went a step farther and took up its habitation within the mussel itself.

*Food and feeding:*—An examination of the stomach and intestinal contents of the various species of mussels of the lake showed no noticeable differences between the food of the different species. Enough of the bottom mud is generally present to give the food mass the color of the bottom on which the mussels are found; thus the stomach-contents of the mussels found in the black bottom of Lost Lake was usually blackish, while that of those found in the lighter bottom at Long Point was grayish. Intermixed, however, with the whole mass was always enough algæ to give it a somewhat greenish tinge, this green being usually intermixed more or less in the form of flakes. A striking contrast between the stomach contents of mussels inhabiting lakes and those found in rivers is the much greater preponderance of organic matter in the food of the lake mussels. The stomach contents of river-mussels is generally chiefly mud, with a few diatoms, desmids, *Scenedesmus* and *Pediastrum* intermixed, as said above. Those of

the  
or  
cid  
(3  
the  
the  
tor  
ab  
va  
sic  
wa  
no.  
dia  
Cy  
Co  
sic  
Pe  
wh  
mu  
en  
th  
me  
fo  
  
th  
ex  
qu  
su  
ap  
in  
na  
ro  
in  
m  
th  
in  
m  
m  
be  
ur  
wi  
of  
at

the lake mussels are almost always full enough of algæ to be more or less flecked with green and sometimes the whole mass is decidedly greenish. On being placed in a vial of preserving fluid (3% formalin was generally used) and shaken, the material from the river mussels always retains the uniform appearance of mud; that from the lake mussels separates, the mud settling to the bottom and the organic material settling as a light flocculent mass above the more solid portion. This top layer is composed of the various plankton elements of the lake, and was found to vary considerably in different lakes. In the Lake Maxinkuckee mussels it was found to consist chiefly of such species as *Microcystis aeruginosa*, *Botryococcus braunii*, *Coelosphaerium kuetzingianum*, various diatoms, such as species of *Navicula*, *Rhoicosphenia*, *Gomphonema*, *Cyclotella*, and *Cocconema*, various forms of desmids, especially *Cosmarium* and *Staurastrum*, various forms of *Scenedesmus*, considerable *Peridinium tabulatum*, and short filaments of *Lyngbya*. *Pediastrum*, both *boryanum* and *duplex* are here, as almost everywhere, rather common objects encountered in the intestines of mussels. Casts of the rotifer *Anuraea cochlearis*, and of the small entomostracan Chydorus, were occasionally encountered. In one of the Lost Lake mussels, *Dinobryon*, an exceedingly frequent element of the mussel-food in Lake Amelia, Minn., but rare here, was found.

No opportunities were had to study the stomach contents during the winter, the mussel work having not been taken up to any great extent during the earlier part of the survey. Mussels obtained quite late in the autumn contained much the same material as in summer. The open and apparently active inhalent and exhalent apertures noted throughout the winter in some individuals would indicate that the mussels—at least some of them—do not hibernate, but carry on life processes more or less actively the year round. The presence of pretty well marked growth-rings would indicate, however, annual rest periods. As diatoms appear to be much more abundant in the water during the winter, it is probable that they enter more plentifully into the mussel's bill-of-fare during the late autumn, winter and early spring than during the summer. In consideration of the mussels as feeders on plankton elements, it is worth while to investigate whether these are not of benefit to the lake as the reducers of excessive amounts of such undesirable elements as *Lyngbya*, *Anabaena* and *Microcystis*, and whether a considerable increase in the mussel population by means of artificial propagation would not clear up the lake to a considerable extent. The following studies of stomach contents and table

of mussel food are by no means exhaustive, but represent hurried examinations and a record of the more easily recognized forms out of a mass of doubtful material. They are intended to be simply suggestive.

Closely connected with the question of food and nutrition is that of the size of mussels. A marked feature of the mussels of Lake Maxinkuckee as well as of the neighboring lakes, is the dwarfing of many of the species, and this is rather difficult to explain, when one considers the large amount of organic material they ingest. The mussels of a few northern lakes examined were thick-shelled and large, so this dwarfness may not be necessarily associated with lake conditions, that is, absence of current. A possible explanation is that of close inbreeding, there being no admixture of new blood with other distant colonies, such as is possible where the lake is in close connection with a large river and its mussel beds.

*Breeding habits, reproduction, etc.*:—The reference to inbreeding above leads to a consideration of breeding and breeding habits. At first glance it would appear that lakes, having no or only feeble currents would make fertilization of the ova of the female mussels largely a question of chance. It is not possible, with the data at hand, to make precise comparisons between number of gravid females of the mussels of lakes and rivers during the proper seasons; but the general impression gained from having examined the mussels of numerous lakes and rivers through the different seasons is that there are fewer of the mussels of the lake that succeed in having their ova fertilized. Gravid mussels are indeed not rare in the lake at the proper seasons, but they seem to be much fewer than one might expect. On October 17, 1907, for example, of 252 *Lampsilis luteola* examined, 41 were of the characteristic female form but only 25 were gravid. Likewise of 18 *Anodonta* examined on the same date, only 2 were gravid. This is a considerably lower percentage than one would expect in rivers at that date. There are other indications that the functions of reproduction are much less prominent in the lake than in rivers. In the height of the spawning season certain species of mussels, especially *Lampsilis ventricosa* and *L. multiradiata*, exhibit, in the neighboring rivers, a very striking appearance, due to the excessive development and high coloration of portions of the mantle near the inhalent aperture. Though both these species are found in the lake, none was observed in this condition. In some rivers in densely crowded beds, moreover, one frequently encounters precocious individuals; small shells, usually apparently only two or three years old but gravid and with the characteristic female contour markedly de-



veloped. This is possibly related to opportunities of fertilization of ova, and is most frequently observed in *L. ventricosa* and *L. luteola*; no such precociously developed mussels were found in the lakes.

A large and well developed female *Lampsilis ventricosa* was transplanted from Yellow River into Lake Maxinkuckee. On being examined two years later in the autumn when this species is usually gravid, it was found to be sterile.

The natural infection of fishes of the lake with the glochidia of the mussels does not appear to be common. The gills of an immense number of fishes were examined for parasites, but no glochidia were noted. Some young bluegill and redeye, exposed to glochidia of *L. luteola* in the autumn of 1912, took very readily.

Very young mussels were either few, or very difficult to find. Diligent search was made for them, especially in the sandy bottom near Long Point, the sand being scooped up and sieved through fine-meshed sieves. Numerous and varied forms of life were thus obtained, such as *Sphaerium*, *Pisidium*, caddis-cases, etc., and rather small but by no means minute examples of *L. luteola*. These young shells were remarkably brightly rayed. Half-grown *Q. rubiginosa* were fairly common in the beds of Lost Lake.

*Proportion of various species in the lake:*—Of a collection of 340 living mussels collected October 17, 1907, at Long Point, 252 were *Lampsilis luteola*, 41 *L. ventricosa*, 21 *Unio gibbosus*, 18 *Anodonta grandis footiana*, 5 *Strophitus edentulus* and 3 *Lampsilis subrostrata*. In deep water *U. gibbosus* and *Anodonta* would have given a higher percentage, and in the Lost Lake beds *Quadrala rubiginosa* would be present in considerable relative abundance.

*Parasites, enemies, and diseases:*—As a general rule the mussels of lakes, ponds and bayous are more heavily infested with parasites than those of swiftly flowing rivers, the probable reason being that in still waters the parasites can migrate more easily from one mussel to another than where there is a swift current. The mussels of the lake are not nearly so badly parasitized as those of the sloughs of the Mississippi, the dead water in the Maumee above the dams, or those of Twin Lakes a few miles to the north. The parasites will be taken up more fully in consideration of the various species of mussels. *Cotylaspis insignis* and several species of *Atax* are the most common parasites. Unlike the mussels of most of our rivers, the mussels of the lakes are comparatively exempt from the ravages of man. A few are killed and used for bait, and now and then a mild case of pearl fever ap-

pears at the lake but is soon cured by the examination of a bushel or two of mussels. On September 22, 1907, a man was seen at the south end of the lake with about a peck of shells which he had opened in a vain search for pearls; on October 8 of the same year a pile of about a half bushel of shells, which had evidently been opened by pearl-ers, was found in Overmyer's woods. Another pearler was seen in 1907 who had collected a few slugs of almost no value. One of the citizens of Culver, in 1906, submitted a small vial of lake baroques for valuation, but they had no worth whatever. The greatest enemy of the lake mussels is the muskrat, and its depredations are for the most part confined to mussels near shore. The muskrat does not usually begin its mussel diet until rather late autumn, when much of the succulent vegetation upon which it feeds has been cut down by frost. Some autumns, however, they begin much earlier than others; a scarcity of vegetation or an abundance of old muskrats may have much to do with this. The rodent usually chooses for its feeding grounds some object projecting out above the water, such as a pier or the top of a fallen tree. Near or under such objects one occasionally finds large piles of shells. The muskrat apparently has no especial preference for one species of mussel above another, but naturally subsists most freely on the most abundant species. These shell piles are excellent places to search for the rarer shells of the lake.

On September 24, 1907, about a bushel of shells, recently cleaned out by muskrats, was found at Long Point where a pier had been removed not long before. The shells were all of rather small size and were in about 18 inches of water. About half were taken and examined. Of these shells 358 were *Lampsilis luteola*, 167 *Unio gibbosus*, 6 *Lampsilis iris*, and 1 *Lampsilis multiradiata*. In the autumn of 1913 freshly opened shells of *Lampsilis glans* were common along shore at Long Point. The first shells killed are rather small and are probably killed by young muskrats.

In the winter after the lake is frozen, great cracks through the ice extend out from shore in various directions, and this enables the muskrat to extend his depredations some distance from shore in definite limited directions. During the winter of 1904 a muskrat was observed feeding on mussels along the broad ice-crack that extended from the end of Long Point northeastward across the lake. The muskrat was about 500 feet from shore. It repeatedly dived from the edge of the ice-crack, and reappeared with a mussel in its mouth. Upon reaching the surface with its catch, it sat down on its haunches at the edge of the creek, and, holding the mussel in its front feet, pried the valves apart with its teeth and

sc  
n  
le  
-al  
ki  
g  
w

wo  
w  
Ti  
re  
lal  
th  
al  
19  
Cu  
At  
Ti  
br  
sa

ert  
fou  
cor  
all  
jet  
ter  
the  
Lal  
gra  
are  
bur  
are  
bea  
of

scooped or licked out the contents of the shell. Some of the larger mussels were too strong for it to open, and a part of these were left lying on the ice. The bottom of the lake near Long Point, and also over by Norris's, is well paved with shells that have been killed by muskrats. Muskrats do not seem to relish the gills of gravid mussels; these parts are occasionally found untouched where the animal had been feeding.

LIST OF SPECIES

1. FLAT NIGGERHEAD

QUADRULA COCCINEA (Conrad)

Rare at the lake; this is a river rather than a lake shell and would be expected in abundance only in fluviatile lakes, or lakes with broad short outlets and vital connection with river faunas. The few living mussels of this species found in the lake probably represent a vanishing remnant of a fauna introduced when the lake had a broader outlet than at present and communication with the river below was more active. A few dead shells were found along the north shore of the lake at various times. On October 25, 1907, a shell 1.75 inches long was found near the railroad bridge at Culver, and in 1909 another small shell was found on the shore at Aubeenaubee Bay. Some fine large examples brought up from the Tippecanoe were planted in the Thoroughfare below the railroad bridge, but they have probably been covered and suffocated by sand.

2. WABASH PIG-TOE

QUADRULA RUBIGINOSA (Lea)

More common in Lake Maxinkuckee than *Q. coccinea*, but nevertheless rather rare, only a few dwarfed shells having been found. In Lost Lake below the Bardsley cottage it was a fairly common species. None of the shells found was of large size, but all were well-formed and handsome. The older shells are almost jet black and peculiarly elongate, with the umbones markedly anterior in position. They look considerably unlike those of either the Tippecanoe or Yellow River, but a form much like the Lost Lake shells was found in the lower course of the Kankakee. No gravid examples were found in the lake. Half grown examples are rather common in Lost Lake beds, but as they are usually buried considerably deeper in the sand than the older shells, they are harder to find. These half-grown shells are of a peculiarly beautiful golden yellow color with a satiny epidermis, and are of the same shape as those found in the neighboring rivers—

that is, the normal or usual shape of the species. The peculiar elongate form of the adult is therefore evidently the product of local influences. The young shells are very iridescent and translucent—much more so than those found in rivers.

*Q. rubiginosa* is at its best a very fair button shell, but the lake shells are too small to work up well. This species appears to be rather rare in lakes. The only lake examples of this species with which the Lost Lake shells were compared were some obtained in Lake Erie. The Lake Erie shells were much more dwarfed, but very solid.

*Food*:—The following is the result of an examination of the material found in the intestines of *Q. rubiginosa* from Lost Lake:

Sample 1. August 2, 1908. Mass fine flocculent rather brownish green material, cohering somewhat in cylinders. Looks as if chiefly organic; not gritty to touch. Organisms present: *Scenedesmus*, *Fragilaria*, *Tetraëdron*, *Navicula*, *Peridinium tabulatum*, *Anuraea*, and *Botryococcus braunii*.

Sample 2. August 20, 1908. A large amount of material; appearance in vial, bottom black, top a fine flocculent sediment. In the top material are *Tetraëdron*, *Scenedesmus*, *Microcystis aeruginosa*, and many disassociated minute cells. Black bottom composed of *Anuraea*, *Lyngbya aestuarii*, a long filament; *Scenedesmus*, many, *Peridinium tabulatum*, *Tetraëdron*, *Epithemia turgida*, *Merismopedia*, cast of *Cyclops*, *Melosira crenulata*, *Gloeocapsa*, *Staurastrum*, *Pediastrum boryanum*, *Gomphonema*, *Chaetophora*, *Cosmarium*, sponge spicule, *Gomphosphaeria aponina*, and *Botryococcus braunii*.

Sample 3. August 20, 1908. A small amount of flocculent brownish material.

*Microcystis aeruginosa*, *Peridinium tabulatum* many and a good many empty cuirasses, *Chydorus*, *Eudorina* a few, *Scenedesmus* common; Diatoms, *Pediastrum duplex*.

Sample 4. August 20, 1908. Fine blue-green flocculent material. *Lyngbya aestuarii*, several filaments; *Microcystis aeruginosa*, common; *Coelosphaerium kuetzingianum*, *Peridinium tabulatum*, very abundant; *Chydorus*, *Anuraea*, *Botryococcus braunii*, *Coelastrum*, *Staurastrum* 1, small, *Navicula*, several.

Sample 5. August 20, 1908. Fine bluish-green material. *Peridinium tabulatum*, abundant; *Cymbella cymbiformis*, *Navicula*, a few; *Anuraea cochlearis*, *Microcystis aeruginosa*, *Chydorus* 1 entire, and other fragments; *Pediastrum duplex*, *Coelosphaerium kuetzingianum*; *Cosmarium*, *Coscinodiscus*, *Scenedesmus*, very common; *Merismopedia glauca*.

Sample 6. August 20, 1908. A small amount of flocculent grayish material.

*Peridinium tabulatum*, abundant, agglutinated in masses; *Microcystis aeruginosa*, very common; *Navicula*, *Staurastrum*, *Cosmarium*, several; *Chydorus*, fragment; *Scenedesmus*, small forms, common; *Pediastrum boryanum*, *Cymbella cymbiformis*, *Tetraëdron*, common; various diatoms; Rotifer, an elongate species; *Merismopedia glauca*; *Coelastrum*, desmids.

Sample 7. August 21, 1908. A small amount of rather coherent fine flocculent greenish material.

*Peridinium tabulatum*, very common; *Anuraea cochlearis*, a few; *Microcystis aeruginosa*, frequent; *Lyngbya aestuarii*, short filament; *Pediastrum boryanum*, diatoms (*Cymbella cymbiformis*; *Cymatopleura*; *Epithemia argus*, *Gomphonema*, *Synedra*), *Tetraëdron*, *Scenedesmus*, occasional; *Dinobryon*, *Staurastrum*, rather slender form.

Sample 8. August 20, 1908. A small amount of flocculent bluish-gray material.

*Peridinium tabulatum*, most abundant; *Coelosphaerium kuetzingianum*; *Pediastrum duplex*; *Microcystis aeruginosa*, *Anuraea cochlearis*, sponge spicule, diatoms (*Navicula*, *Cymbella*, etc.), *Scenedesmus*.

Sample 9. August 20, 1908; a fair amount of flocculent grayish-brown material with a greenish cast.

*Peridinium tabulatum*, most abundant; *Microcystis aeruginosa*, *Anuraea cochlearis*, *Staurastrum*, *Pediastrum duplex*, *Botryococcus braunii*; *Tetraëdron minimum*, *Coelosphaerium kuetzingianum*; *Pediastrum boryanum*, *Chydorus*, *Lyngbya aestuarii*, *Gloeocapsa*, diatoms—*Cymbella cymbiformis*, *Navicula*.

### 3. SPIKE

#### UNIO GIBBOSUS Barnes

This mussel, known among clammers as the "spike" or "lady-finger" is, next to *Lampsilis luteola*, the most abundant shell in the lake. It is found wherever the other mussels are; that is, in sandy or somewhat marly bottom in rather shallow water most of the way around the lake, and in the shell-bed in Lost Lake below Bardsley's. In Lake Maxinkuckee one of the best beds is at Long Point. It is abundant also at Norris Inlet, and by McDonald's and Farrar's.

No very young of this species were found in the lake; they are, however, hard to find in numbers anywhere, even in rivers where

the species is abundant, except in cases where portions of the river go almost dry, and this of course never happens to the beds in the lake. The half-grown examples are solid, rather cylindrical shells, the same neat form that is known as the "spike" among the clammers. The old shells develop into a peculiar form, being flattened, arcuate along the ventral border and very thin posteriorly, so that they usually crack badly in drying; they represent the form described by Simpson as var. *delicata*. In general outline they remind one somewhat of *Margaritana monodonta*. This form is not strictly confined to the lake; some similar shells were collected in the Wabash near Terre Haute.

As found in the lake, *Unio gibbosus* is very constant in its characters, the only noteworthy difference between individuals being the change in shape already referred to as being due to age. In rivers this shell exhibits considerable variation in shape, size, color of nacre, etc., but the shells of the lake are quite constant in almost every respect. The nacre is a deep purple, never varying to pink or white as it frequently does in rivers.

Like *Lampsilis luteola* this species is frequently preyed upon by muskrats and the cleaned out shells are common where these rodents have had their feasts.

Although *U. gibbosus* of the Tippecanoe River near the mouth of the Outlet is very commonly infested with a distomid parasite along the hinge-line which brings about the formation of irregular baroques, this parasite does not occur in the lake so far as known. Small species of *Atax* are common parasites of this species in the lake, and in 1909 one was found affected by the large *Atax ingens*.

Even the large strong river shells of *Unio gibbosus* have no value in the manufacture of button shells because of their purple color and lack of luster. (The white-nacred shells are sometimes used.)

The only other lake examples with which the Lake Maxinkuckee specimens of this species have been compared, are some collected in Lake Erie at Put-in-Bay. The Lake Erie shells are much unlike the Maxinkuckee specimens, being short, humped and remarkably solid and heavy. Similar shells to those of Lake Erie are found in some of the small southern rivers.

We have no notes referring to gravid examples in the lake. This was probably because the most active work in collecting and examining mussels was carried on in the autumn, and the breeding period of this species is in early summer.

## 4. ALASMDONTA CALCEOLUS (Lea)

Judging from the dead shells found scattered along shore, this is not a particularly rare species in the lake. The shells were found most abundantly along the north shore of the lake, although they were also found along the east and southeast portion and were not infrequent between Arlington and Long Point. No living examples were found. On account of its small size and its habits this is a rather difficult species to find, even where common, except under favorable conditions such as exceptionally low water, when the mussels move about more or less. Nothing was therefore learned of its habits in the lake. In the Tippecanoe River near De-long, Ind., this species was rather common in stiff blue clay near shore, and it is fairly abundant in Yellow River at Plymouth. Here, although the dead shells were common, the living examples were difficult to find until, during a period of very low water, they began actively moving about and could be tracked down. The species, which reaches an unusually large size in Yellow River, was there found gravid in autumn (September and October). The glochidia are of the Anodonta type, chestnut-shaped or rounded-triangular in outline, with large hooks at the ventral tips of the valves.

## 5. FLOATER

## ANODONTA GRANDIS FOOTIANA Lea

Although the genus *Anodonta* is generally regarded as the "Pond-mussel" *par excellence*, the species of which might naturally be expected to be at home in lakes and ponds and thrive in such places even better than in rivers, the Anodontas of Lake Maxinkuckee show, along with the river-species proper, the dwarfing influence of the lake. Moreover, *Anodonta* is not, as one might naturally expect, the most abundant mussel in the lake, but is outnumbered in some beds at least, by both *Lampsilis luteola* and *Unio gibbosus*. Its relative scarcity in some of the shore beds is in part made up by its wider distribution in the deeper waters of the lake than the others reach, and on its presence on the isolated bars, where it was occasionally taken up by the dredge.

On account of the great variability of *Anodonta grandis* and the difficulty in distinguishing the various forms, particular attention was paid to this species as found in the lake, and the lake specimens were compared with numerous examples from the neighboring lakes and river. No Anodontas were found in the Tippecanoe River near Lake Maxinkuckee Outlet, and we were therefore

unable to compare our lake specimens with the form that would be most interesting in this connection.

The mussels of Tippecanoe Lake at the head of Tippecanoe River were examined in this connection. Blatchley (Indiana Geological Report for 1900, p. 190) has reported *Anodonta grandis* as common, and the subspecies *footiana* as frequent in Tippecanoe Lake. The Anodontas of that lake differ markedly both in the size and shape of the individuals from those of Lake Maxinkuckee. The difference in size can be easily explained by the more favorable conditions in Tippecanoe Lake. This body of water is more fluviatile than Lake Maxinkuckee, being directly connected with the Tippecanoe River, which is already a fairly large stream when it leaves the lake, and the mussel beds of the lake and river are continuous. The upper part of Tippecanoe Lake is exceptionally favorable for Anodontas; the living mussels are large and abundant, and the dead shells almost pave the bottom near shore, several dead shells often being telescoped within each other. Some of the shells reached a size not often surpassed in the neighboring rivers, one example measuring 172.5 mm. long, 95 mm. high and 65 mm. in diameter. A few were thickened with a tendency to form half pearls, but most were thin. A number of the shells approached *Anodonta corpulenta* in general form, and one flattened, rounded shell resembled *A. suborbiculata*. The Anodontas from other lakes of the Tippecanoe River system, such as Center Lake and Eagle Lake near Warsaw, resembled those of Lake Maxinkuckee, but were generally smaller and shorter.

The Anodontas of Lake Maxinkuckee were also compared with those of Yellow River a few miles to the north, and with the various lakes of the Kankakee system, including Upper Fish Lake, Lake of the Woods (Marshall Co), Pretty Lake, Twin Lakes, Bass Lake and Cedar Lake. Some of the Yellow River Anodontas were normal, oval shells such as are common in the rivers of northern Indiana; a few were exceptionally thin and exceedingly inflated, resembling *A. corpulenta*. Those of Upper Fish Lake—originally a fluviatile lake containing other fluviatile shells such as *Q. undulata*—were large shells like those of Tippecanoe Lake. The Anodontas of each of the other lakes differed more or less from those of the others, though all probably had a common origin. The only lake of this group, the Anodontas of which closely resembled those of Lake Maxinkuckee, is Bass Lake, and even here the shells were somewhat different, being smaller and with the epidermis more deeply stained. Even the Anodontas of Lost Lake differ slightly from those of Lake Maxinkuckee, being somewhat more inflated



and with the epidermis rather green than brown, and in having the shell usually somewhat thinner. Some of the shells near the outlet of Lost Lake are exceedingly thin, some of them so much so that ordinary print can easily be read through them; they are so fragile that it is almost impossible to keep them.

Of the collection from Lake Maxinkuckee, mostly from Long Point, 26 examples were carefully compared. The smallest measured 68 mm. long, 38 mm. high and 24.6 mm. in diameter, and the largest 93.5 mm. long, 50 mm. high and 37 mm. in diameter. Among variant forms was one female, gravid when collected, which was unusually elongate, its measurements being 86 mm. long, 43.5 mm. high and 32.5 mm. in diameter. In outline this shell closely resembled *Anodontoides ferussacianus subcylindraceus*.

Some of the larger specimens are rather humped and arcuate, the ventral margin of one being somewhat concave. This is a variation which is quite likely to occur in old shells of any species.

Although gravid Anodontas were found rather frequently during the late autumn, no infected fishes were seen, and no young were found.

The Anodontas of the lake are fairly free from parasites, a few *Atax* and *Cotylaspis* and occasionally a few distomids on the mantle next to the umbonal cavity being the only ones present in any numbers. In some of the other lakes the Anodontas were very badly infested; a colony found in one of the Twin Lakes being infested to a remarkable degree by a distomid which formed cysts in the margin of the mantle.

*Food and Parasites of various examples:*—The following is the result of the examination of various examples of Anodontas from the lake.

Sample 10. Vial containing intestinal contents of *Anodonta grandis footiana*, Lost Lake, Sept. 7, 1908. The vial contains a considerable amount of material (in formalin) which was separated into black fine mud below and fine flocculent light green above. Upper portion—*Microcystis aeruginosa* most common; *Peridinium tabulatum* some; *Pediastrum boryanum*; *Melosira crenulata*, a few filaments; *Coelastrum microporum*, *Botryococcus braunii* and *Scenedesmus*. Bottom layer—*Lynbya aestuarii*, *Microcystis aeruginosa* very common; *Peridinium tabulatum*, *Anuraea cochlearis*, *Cymbella cymbiformis* and *Navicula*.

Sample 11. Food of *Anodonta grandis footiana*, Lake Maxinkuckee, near Norris Inlet, Aug. 20, 1908. A good mass of flocculent fine green material; no mud.

*Microcystis aeruginosa* most common, *Melosira* filament, *Oscillatoria*, short filament; *Anuraea cochlearis* several; *Cymbella cymbiformis*; *Gomphosphaeria aponina*; *Peridinium tabulatum*; *Coclosphaerium kuetzingianum*, *Lyngbya aestuarii*, *Epithemia argus*, *Chydorus*, and what appears to be fragments of *Ceratium hirundinella*.

Sample 12. *Anodonta grandis footiana*, near Norris Inlet, Lake Maxinkuckee, Aug. 20, 1908; a small mass of flocculent blue material.

*Microcystis aeruginosa* most abundant; *Lyngbya aestuarii*, *Melosira*, *Epithemia*, *Anuraea cochlearis*, *Pediastrum boryanum*, *Cosmarium intermedium* and a few others, *Staurastrum*, *Spirulina* and *Pediastrum duplex*.

Sample 13. *Anodonta grandis footiana*, 97 mm. long. Edge of Lake Maxinkuckee east of Norris Inlet, Aug. 29, 1908.

Parasites, 9 *Atax*, free among gills. Mussels gravid, with anterior end of shell indented and with some brown spots on the nacre. Food mass fine golden brown, abundant in quantity, containing *Anuraea cochlearis* many; *Microcystis aeruginosa* most abundant element; *Lyngbya aestuarii* frequent; *Scenedesmus*, a few; *Botryococcus braunii* frequent, *Cymbella cymbiformis*; *Staurastrum*, *Navicula*; *Fragilaria*; *Chydorus*, a few; *Coelosphaerium kuetzingianum*; the diatoms are not abundant.

Sample 14. *Anodonta grandis footiana* apparently old, 90 mm. long, near Norris Inlet, Lake Maxinkuckee, Ind., Aug. 29, 1908, the shell stained somewhat brown inside, with one steel-blue stain on the right valve anteriorly.

Parasites; *Atax* 7, large, full of eggs, one small, one very small, these all free among the gills; *Cotylaspis insignis* 1, in axil of gill.

Food abundant; *Microcystis aeruginosa* abundant, *Lyngbya aestuarii* common, *Pediastrum duplex*, *Botryococcus braunii*, a few; *Cosmarium*; *Anuraea cochlearis* several; *Scenedesmus*; *Ankistrodesmus*, and many diatoms, among which are *Cocconeis pediculus*, *Melosira*, *Gomphonema*, *Navicula*, *Epithemia turgida*, etc.

Sample 15. *Anodonta grandis footiana*, 101 mm. long, Lake Maxinkuckee, near shore, by Norris Inlet, Aug. 29, 1908.

Parasites, 5 *Atax*, free in gills, some full of eggs, 2 smaller in size, larval *Atax* (black) scattered in gills. *Cotylaspis insignis* 2, axil of inner gill.

A large amount of food material in intestines, very fine, of a yellowish-brown color.

*Microcystis aeruginosa*, *Anuraea cochlearis*, *Lyngbya aestuarii*, *Botryococcus braunii*, *Coelosphaerium kuetzingianum*, *Cosmarium*,

Ne  
di  
  
bc  
M  
sc  
  
M  
ab  
an  
Sc  
lar  
sr  
  
bc  
M  
m  
m  
oc  
E  
M  
  
M  
Pe  
yo  
  
fr  
fil  
Pe  
Ne  
se  
  
ne  
dis  
  
ae  
Ne  
  
tra  
30  
in  
sid

*Navicula*, an elongate form, *Cymbella cymbiformis*, *Pediastrum duplex*, *P. boryanum*; red cysts apparently of *Peridinium*.

Sample 16. *Anodonta grandis footiana*, 90 mm. long, sandy bottom of Lake Maxinkuckee near Norris Inlet. Aug. 29, 1908. Mussel gravid. Parasites, *Atax*, 3, free among gills, *Atax* embryos scattered through gills.

Food material scarce, fine golden brown in mass, consisting of *Microcystis aeruginosa* abundant; *Coelosphaerium kuetzingianum* abundant, *Lyngbya aestuarii*, a few filaments; *Anuraea cochlearis* and another rotifer; *Botryococcus braunii*; *Sorastrum*, *Coelastrum*, *Scenedesmus*, *Pediastrum duplex*, *Navicula* several, *Melosira tabulata*, *Synedra*, *Epithemia turgida*, *Cymbella cymbiformis*; other small diatoms rather numerous; *Cosmarium*, a few.

Sample 17. *Anodonta grandis footiana*, 93 mm. long. Sandy bottom of Lake Maxinkuckee near Norris Inlet, Aug. 28, 1908. Mussel gravid. Parasites, 1 *Atax*, free among gills. Intestines almost empty. *Microcystis aeruginosa* one of most abundant elements, *Lyngbya aestuarii*, *Coelosphaerium kuetzingianum*, *Botryococcus braunii*; *Cosmarium*, *Pediastrum*, *Cocconeis pediculus*, *Epithemia turgida*; *Navicula*, (1 actively moving) *Gomphonema*, *Melosira tabulata*, *Anuraea cochlearis*, *Chydorus*.

Sample 18. *Anodonta grandis footiana*, 95 mm. long, Lake Maxinkuckee near Norris Inlet, Aug. 29, 1908. Mussel gravid. Parasites, 6 *Atax* free among gills, one a minute red species. Many young *Atax* embryos in inner side of mantle, not in gills.

Food material golden brown with some green intermixed, very fine. *Microcystis aeruginosa*, common; *Lyngbya aestuarii*, a few filaments; *Coelosphaerium kuetzingianum*; *Botryococcus braunii*; *Pediastrum duplex*; *Anuraea cochlearis* a few; *Epithemia turgida*; *Navicula*, common; *Cymbella cymbiformis*; *Cocconeis pediculus*, several; *Cosmarium*; *Chydorus*.

Sample 19. *Anodonta grandis footiana*, Lake Maxinkuckee, near Winfield's. Mussel gravid. Parasites, young *Atax* in gills; distomids on mantle; (a slug pearl near hinge.)

Food: *Botryococcus braunii*; *Microcystis aeruginosa*; *Lyngbya aestuarii*, *Coelosphaerium kuetzingianum*, *Pediastrum duplex*, *Navicula*, *Cymbella cymbiformis*.

Sample 20. *Anodonta grandis footiana*, Lost Lake. Young transparent shell, gravid; length 77 mm., height 41 mm., diameter 30 mm.; live weight 1 oz., shell  $\frac{1}{4}$  oz. Parasites, several *Cotylaspis insignis* in axil of gills. Food chiefly *Microcystis aeruginosa*; considerable *Botryococcus braunii*.

Sample 21. *Anodonta grandis footiana*, Lost Lake. Parasites,

1 young *Atax* in gill; *Cotylaspis insignis* in axil of gill. Food chiefly *Microcystis aeruginosa*, a little *Botryococcus braunii*, *Lyngbya aestuarii* and *Pediastrum boryanum*.

6. PAPER-SHELL

*ANODONTA IMBECILLIS* Say

A single specimen.

7. SQUAWFOOT

*STROPHITUS EDENTULUS* (Say)

Not very common in the lake. Occasional shells can be picked up along shore, especially between Long Point and Arlington, and along the north shore. Living examples were also taken in small numbers from the mussel bed at the mouth of Norris Inlet, and at Long Point. In a collection of about 300 living mussels collected at the latter place in the autumn of 1907, only 3 were of this species.

As found in the various rivers of the country, this is one of the most variable of shells, and the exact limits of the species and its various forms are not yet well worked out. The lake examples, though differing considerably from those of the neighboring rivers and from river shells in general, do not exhibit a very large range of variation. They are all markedly dwarfed, the average length being about 2½ inches or 63.5 mm. long. All have a well-developed rounded posterior ridge. The epidermis is deeply stained, that of the exposed portion of the shell being a rich yellowish brown, while the anterior portion—in the living shell buried in the soil of the bottom—is a deep, shining, brown black. The anterior margin is not nearly so heavy and produced as one frequently finds it in river examples. The beaks of the lake shells are not so angular as they usually are in river shells, and the high wavy ridges are more numerous and pronounced. In the Maxinkuckee shells, also, a number of fine hair-like lines or ridges, much like growth lines, extend along the posterior border of the umbone, parallel with the posterior ridge of the earlier stages of the shell.

The nacre of the lake shells is a rich rosy salmon. Unlike the salmon color of "*Anodonta salmonia*" this is a natural color, not due to diseased conditions; the nacre surface is very smooth and the color extends deeply into the shell. In some cases the inner nacreous surface appears to be a secondary thickening of the shell, laid on the older portions like an enamel. Below this extra nacreous deposit the growth lines are very distinct on the

in  
lin  
up  
the  
mi  
in  
co  
  
est  
  
alc  
sh:  
ab  
an  
fa  
fev  
in  
alc  
mu  
lat  
  
spe  
on  
of  
in  
sm  
  
sca  
is  
off  
at  
  
riv  
she  
anc  
ray  
anc

inner surface of the shell. The rest periods are distinct black lines, often plainly visible through the translucent shell when held up to the light. Rays are always invisible by reflected light in the lake shells, but in some examples they were visible by transmitted light. The animal has orange-colored flesh. The few living examples examined indicate that parasites are common: one contained three old *Atax ypsilophorus*, and several young.

One gravid example was found, October 17, 1907. The youngest example found was 42 mm. long and exhibited four rest periods.

8. LAMPSILIS GLANS (Lea)

Fairly common in the main lake; dead shells are often found along shore, and occasionally the living mussels are to be seen in shallow water at the various mussel beds in the lake. It is quite abundant along the edges of the thoroughfare joining the lakes, and is common in Lost Lake. The examples found in the thoroughfare and Lost Lake were of unusually large size; this is one of the few species of mussels which are as large or larger in the lake than in the neighboring rivers. *L. glans* appears to prefer shallow water along shore. A good number of shells recently cleaned out by muskrats was found near the water's edge at Long Point in the late autumn of 1913.

In the Tippecanoe River at Delong this was a very abundant species in the greasy whitish blue clay along shore, and was here one of the favorite morsels of the muskrat. With the exception of *Micromya fabalis* this is the smallest species of mussel found in the lake. It can be easily recognized by its black epidermis, small size and purple nacre.

9. RAINBOW-SHELL

LAMPSILIS IRIS (Lea)

Rather common in the lake in shallow water near shore, found scattered among the other species in the various shell-beds. There is a good colony in the Lost Lake bed, and it is fairly abundant off the depot grounds, by Kruetzberger's pier, at Long Point, and at the bed near the mouth of Norris Inlet.

The lake shells differ markedly from those of the neighboring rivers, so much so that it is easy to separate the lake and river shells at a glance. The lake shells are considerably more elongate, and the epidermis is stained a deep brown, mostly concealing the rays; when these are visible they are brownish rather than green, and the umbones are rather eroded. The shells, indeed, resemble

somewhat the males of *L. subrostrata*, with which they are associated. The lake shells exhibit a tendency to have their posterior margin somewhat broader than that of the river shells, and the shells are flatter at the posterior tip, becoming somewhat produced. The river shells are more solid and heavy.

*Lampsilis iris* is one of the few species of mussels which does not show a marked decrease of size in the lake; indeed, some of the larger lake examples run actually larger than those from the neighboring rivers. Some of the largest lake shells examined have the following dimensions:

No.	Length mm.	Alt. mm.	Diam. mm.
1	69.6	37.3	21
2	65.9	34.9	21
3	68.0	34.6	22
4	64.9	35.8	22.7
5	67.0	36.8	20.9
6	67.7	33.8	21.5

No young shells were found, even the smallest appear rather old. The smallest three measure:

Length mm.	Alt. mm.	Diam. mm.
41.4	21.2	12.5
38.9	21.5	12.5
37.0	20.0	12.3

For comparison with the lake shells, the dimensions are given of the largest two shells found in Yellow River:

No.	Length mm.	Alt. mm.	Diam. mm.
1	67.0	34.5	22.9
2	64.0	33.5	21.0

Only one gravid example was found; this was obtained at Lost Lake bed Sept. 7, 1908.

Of all the species of mussels in the lake, *L. iris* has the best connection, through scattered individuals along the Outlet, with the shells of the Tippecanoe River, a few shells having been found almost through the whole length of the Outlet. The outlet shells, like those of the rivers, are brightly rayed. The species is abundant in the Tippecanoe River at Delong. A number of examples were noted in spawning condition there in late August and early September in 1908. Observations in the Maumee River would indicate that these species, *L. parva* and *L. multiradiata*, do not have exactly the same breeding season as many other species of *Lampsilis* (*luteola*, *recta*, *ligamentina*, etc.) but are sometimes fertilized in July, spawning in August and September. Being small and

an early developing species, it is probable that they have somewhat different habits: indeed it is possible that they have more breeding seasons per year than the other species.

The Tippecanoe mussels of this species were a favorite food of the muskrat, and were killed in great numbers every autumn, the dead shells being thickly strewn along the bank, or piled in heaps at the bases of rocks which the rodent had used as a feeding place.

*Lampsilis iris* has a well marked tendency in the lakes and Outlet to produce pearls and baroques, but these are too small to be of any value.

10. LAMPSILIS SUBROSTRATA (Say)

*Lampsilis subrostrata* reaches its best development along the muddy shores of lagoons, not being perfectly at home either in swiftly flowing streams or in perfectly quiet lakes, although occasional examples may be found in either. It is considerably more abundant in Lake Tippecanoe and Upper Fish Lake than in any other Indiana lakes examined. Along the edges of the Mississippi sloughs it is fairly common and reaches a large size, often distinguished with difficulty from *Lampsilis fallaciosa* except for the thinness of the shell and the black epidermis. It is rare in Lake Maxinkuckee, only a few examples having been obtained from the mussel bed near Norris Inlet. It is much more common in Lost Lake in the large bed along shore south of the Bardsley cottage. Mr. Blatchley, in a short report on the mollusks of the lake, (25th annual report, Department of Geology and Natural Resources of Indiana, 1900, p. 250), says of this species: "Not common in the main lake; more so in the muck and mud along the margins of Lost Lake, where a well-marked variety, with a larger and broader beak, was taken. A specimen of this was sent, among others, to Mr. Chas. T. Simpson, of the Smithsonian Institution, for verification. In his reply he says: 'The variety of *subrostratus* which you send is, so far as I know, confined to northern Indiana. It is quite remarkable, and would seem to be almost a distinct species. I have seen quite a number of specimens of it, and at first thought it a variety of *U. nasutus*, but there seem to be intermediate forms connecting it with *U. subrostratus*.'"

With the exception of the differences due to sex, all the Maxinkuckee and Lost Lake shells are very uniform in appearance, much more so than *L. luteola*, and are hardly distinguishable from examples from Lake Tippecanoe, Upper Fish Lake, or a specimen collected in the Wabash River at Terre Haute by Dr. J. T. Scovell.

They are dark brown in color with very faint rays. The species appears to be rare in the Tippecanoe River at Delong. One example was obtained there, which is somewhat shorter and stouter than those of the lake, and not so badly stained; it shows faint rays posteriorly. The Lost Lake shells are somewhat larger than those found at the other lakes. No young were found, the smallest shell obtained being a half-grown example. One gravid specimen was found at Lost Lake Sept. 7, 1908. The marsupium closely resembles that of *L. iris*, being a kidney-shaped mass filling the hinder portion of the outer gill, this mass being marked into segments by rather deep radiating furrows. The very edge of the marsupium is white beyond the dusky submarginal area, the white making a chain-like area at the edge of the gill. Like *L. iris*, this species has a tendency to form pearls, but they are too small to be of any value.

*Food of individuals*:—The following is the result of the examination of the contents of the intestines of *L. subrostrata* from Lost Lake at various dates.

Sample 22, August 20, 1908. A small amount of flocculent bluish-gray material.

*Peridinium tabulatum* abundant; *Microcystis aeruginosa* abundant; *Anuraea cochlearis*; *Pediastrum boryanum*; Diatoms—*Synedra*, *Cymbella cymbiformis*.

Sample 23, August 20, 1908. A very small amount of flocculent grayish material.

*Peridinium tabulatum* a few; *Microcystis aeruginosa* a little; *Pediastrum boryanum*; *Cosmarium*, *Tetraëdron minimum*, *Scenedesmus*, *Euglyphia alveolata*; *Peridinium*, a small, sharp-spined form. Diatoms make up the greater part, including *Cymbella cymbiformis*, *Navicula*, *Fragilaria*, *Coscinodiscus*, and *Epithemia*.

Sample 24, September 7. A large amount of material, black mud below, greenish flocculent material above. The upper portion contains chiefly *Botryococcus braunii* and *Microcystis aeruginosa*. Bottom portion—*Microcystis aeruginosa* common; *Botryococcus braunii*; *Peridinium tabulatum*, *Peridinium*, a small spined species; *Scenedesmus*, frequent. *Staurastrum*, *Pediastrum duplex*; *Coelastrum* a few; *Anuraea cochlearis*, *Tetraëdron*, *Docidium*, *Coelosphaerium kuetzingianum*, sponge spicule, *Lyngbya aestuarii*. Diatoms,—*Synedra*, *Navicula*, *Gomphonema*, etc.

I  
ican  
ica  
grea  
I  
four  
the  
adm  
tom  
Bay  
loca  
fror  
Auk  
it w  
tag  
  
tho  
ne  
Lak  
Lak  
spe  
sm  
The  
abc  
ton  
  
La  
int  
  
she  
thi  
the  
un  
Th  
fla  
fo  
sp  
sil  
th  
no



11. FAT MUCKET

LAMPSILIS LUTEOLA (Lamarck)

*Lampsilis luteola* is the most widely distributed of the American Unionidæ, its range extending over nearly all of North America east of the Rocky Mountains. It lives and thrives under a great variety of conditions, being frequent in both lakes and rivers.

In Lake Maxinkuckee this is the most common mussel, being found almost everywhere in water from 2 to 5 or 6 feet deep where the bottom is suitable. It prefers a rather solid bottom with some admixture of sand or gravel, but occurs also even where the bottom is of a rather firm peaty nature as in some places in Outlet Bay. It is, however, rather scarce and widely scattered in such localities. The best beds are found at Long Point, at Farrar's, in front of McDonald's, by the old Kruetzberger pier, and in Aubeenaubee Bay off from the Military Academy. In Lost Lake it was abundant in the large mussel bed below the Bardsley cottage, and a few shells were found in the north end of the lake.

The Lake Maxinkuckee shells are smaller and thinner than those of the rivers; they closely resemble those of most of the neighboring lakes with which they were compared, such as Twin Lakes, Pretty Lake, Bass Lake, etc. The *L. luteola* of Upper Fish Lake are much larger and more like river shells. Compared with specimens of more remote lakes, those of Lake Erie are much smaller, more solid and not stained, the rays being quite distinct. The *L. luteola* of Lake Pokegama, Minn., are unlike any of those above cited, being large, thick and heavy, furnishing excellent button material.

*Lampsilis luteola* is represented in Lake Maxinkuckee and Lost Lake by 2 forms; although these forms are well connected by intergrades the extremes are pretty markedly distinct.

The colony in Lost Lake is composed of compressed, elongate shells, almost as large as those found in rivers, but considerably thinner. It is in the females of this group, and only in part of them, that the greatest variation occurs. The males are not much unlike the ordinary well-known form of the neighboring rivers. The most strongly aberrant females are markedly compressed, and flare out broadly in the post-basal region. The umbones are far forward and they remind one somewhat in contour of the marine species, *Modiola plicatula*. Some of them closely resemble *Lampsilis radiata* of the Atlantic drainage. The Lost Lake mussels of this species are stained a peculiar attractive ash-gray which does not greatly obscure the rays. They are not so heavily encrusted

with marl as are those in the Lake Maxinkuckee beds. Typical Lake Maxinkuckee specimens are dwarfed and stained a deep brown, which obscures the rays. Most of them are thickly coated posteriorly with incrustations of marl. It is principally this species which has associated with it the little water-beetle, *Stenelmis undulatus* Blatchley. At Long Point, where *L. luteola* is the most common mussel, examples of the peculiar Lost Lake form are rather frequent. In comparing sets of shells from the various mussel beds of the lake—Long Point, Farrar's and the Norris Inlet beds—it was noted that the mussels of each bed, as one approached the upper portions of the lake, averaged somewhat smaller.

As regards food, movements, reproduction, etc., *L. luteola* does not differ greatly from the other mussels of the lake with the exception that it appears to be considerably the most active species in the lake. A few were observed moving about during the winter of 1900-1901. The deep water individuals rarely move about at all. In the autumn of 1913 the migration of those near shore into deep water was strikingly shown by a series of numerous furrows, with a mussel at the deep water end and extending from shore outward near Long Point.

As with the other mussels of the lake, reproduction is a rather inconspicuous phenomenon, not attended with the marked display common in the larger river examples. Of 252 examples collected at Long Point Oct. 17, 1907, 25 contained glochidia in the gills, some being very full and much distended. One was found gravid May 24, 1901, and on August 22, 1906, some in Lost Lake appeared to be about ready to spawn.

The young of this species were found rather frequently in the lake, much more frequently, indeed, than any other kind. The smallest examples were obtained while sieving sand for Sphaeriums at Long Point. These young mussels live buried in the fine sand near shore. Specimens up to about a half inch long are very crinkly, being covered with narrow elevated parallel ridges, generally 5 in number, each consisting of 2 open loops placed end to end, the sides of the loops being roughly parallel with the ventral margin of the shell; the ends where they join form a sharp curve upward toward the umbone. These double loops are followed by a number of broken irregular ridges. The markings just described persist on the umbones of the older shells until eroded away. The half grown shells are beautifully rayed with green on a whitish background. As the shells grow older they become gradually stained a deep uniform brown, obscuring the rays.

ab  
anc  
lik  
spe  
  
val  
rou  
pla  
she  
but  
mu  
Mi:  
we  
cor  
ast  
ma  
loa  
  
len  
to  
wh  
kuc  
  
vid  
  
gra  
wei  
you  
of  
bra  
  
leng  
she  
ma  
bra  
  
live  
33  
egg  
aer.

Most of the mussels of the lake are slightly parasitized, none abundantly; they contain a few examples of a small reddish *Atax*, and a few *Cotylaspis insignis*. A small round-worm, somewhat like a vinegar eel, was found very active in the intestine of one specimen; it was probably parasitic.

Small irregular pearls or slugs are produced but they are of no value. In some rivers this species produces an abundance of small round pearls. Some of the pearl-bearing river specimens were planted in the lake in 1912 to see if they would infect the lake shells. The *Lampsilis luteola* of the rivers is a fair button shell, but the Lake Maxinkuckee shells are too small and thin to have much value. It is a remarkable fact that in Lake Pokegama, Minn., *L. luteola* grows abundantly in shallow bottom among the weeds, and there produces a handsome thick heavy shell, one indeed concerning which the pearl button manufacturers are very enthusiastic, so much so that the shells at that distant point from the market brought \$22.00 per ton; in the summer of 1912, two carloads of these shells were shipped to Europe.

Just why the Lake Maxinkuckee shells are not like the excellent ones of Lake Pokegama remains as yet unanswered, but seems to be largely a question of breed. It would certainly be worth while to introduce the Lake Pokegama breed into Lake Maxinkuckee.

Following is the result of the examination of various individuals of the Maxinkuckee and Lost Lake shells:

Sample 25. *L. luteola*. Lost Lake, Sept. 7, 1908. Mussel gravid. Length 100 mm., altitude 62 mm.; diameter 33 mm. Live weight  $3\frac{1}{2}$  oz.; shell  $1\frac{1}{4}$  oz. Parasites—7 free *Atax* among gills, young *Atax* in gills and numerous *Atax* eggs on interior surface of mantle. Food chiefly *Microcystis aeruginosa*, *Botryococcus braunii*, *Lyngbya aestuarii*, *Melosira*, *Navicula*.

Sample 26. *L. luteola*, Lost Lake, Sept. 7, 1908; mussel gravid; length 95 mm., alt., 60 mm., diam., 38 mm.; live weight  $3\frac{3}{4}$  oz.; shell  $1\frac{1}{4}$  oz. Parasites, 7 free *Atax* in gills, and *Atax* eggs in the mantle. Food—chiefly *Microcystis aeruginosa*, also *Botryococcus braunii*, *Navicula*, *Lyngbya aestuarii* and *Anuraea cochlearis*.

Sample 27. *L. luteola*, Lost Lake by Bardsley's, Sept. 7, 1908; live weight  $3\frac{1}{2}$  oz.; shell  $1\frac{1}{2}$  oz.; length 97 mm., alt. 54 mm., diam. 33 mm. Parasites—7 free *Atax* among gills; many small red eggs of *Atax* on inner surface of mantle. Food chiefly *Microcystis aeruginosa*, *Botryococcus braunii* and *Navicula*.

Sample 28. *Lampsilis luteola*, Lost Lake, Sept. 7, 1908. Live

weight  $3\frac{3}{4}$  oz.; length 104 mm., alt. 54 mm., diameter 33 mm. Parasites, *Atax* 6, free among gills, eggs of *Atax* on inner side of mantle, young in pits on side of foot. Food, *Microcystis aeruginosa* most common. *Lyngbya aestuarii*, *Navicula*, *Melosira*, *Anuraea* and *Cymbella*.

Intestinal contents of two examples of *L. luteola* obtained in Lake Maxinkuckee Aug. 27, 1908, near the shore just north of the ice-office gave the following results:

Sample 29. *Microcystis aeruginosa*, main mass; *Anuraea cochlearis* a few; *Botryococcus braunii* rather common; *Cymbella cymbiformis*, one; *Lyngbya aestuarii*, 1 filament; *Navicula*, 2 examples; *Synedra*, a few.

Sample 30. *Microcystis aeruginosa* main mass; *Botryococcus braunii*, very common; *Lyngbya aestuarii*, several filaments; *Anuraea cochlearis* a few; *Synedra* some; *Navicula* one example, very lively; *Cosmarium* one round worm like vinegar eel, very lively.

Sample 31. Lost Lake, 1908. A good mass of material, blackish below, flocculent greenish above.

*Lyngbya aestuarii*, a few filaments; *Microcystis aeruginosa*, quite abundant; *Anuraea cochlearis*; sponge spicule, *Pediastrum duplex*, *Staurastrum*, *Botryococcus braunii*, *Peridinium tabulatum*, a few; *Peridinium*, a small spiny species 1; *Pediastrum boryanum*; several diatoms—*Navicula*, *Coscinodiscus*, *Melosira*, *Cymbella cymbiformis*; *Microcystis* is the most abundant element, *Peridinium* is rather scarce.

Sample 32. Lake Maxinkuckee, Aug. 27, 1908. A small amount of brownish green flocculent material.

*Anuraea cochlearis*, quite frequent; *Lyngbya aestuarii*, short filament; *Peridinium tabulatum*, a few; *Coelastrum microporum*; *Coelosphaerium kuetzingianum*; *Pediastrum boryanum*; *Scenedesmus*, very few; *Chydorus* fragment. Diatoms,—*Epithemia turgida*, *Navicula*, *Cymbella cymbiformis*, *Gomphonema*, *Coscinodiscus*.

Sample 33. Lake Maxinkuckee, Aug. 27, 1908. A fair amount of brownish green material, muddy below, flocculent green above. The green top material consisting chiefly of *Microcystis aeruginosa*; with some *Anuraea cochlearis*; *Lyngbya aestuarii*; *Microcystis aeruginosa*; *Bulbochaete* bristle; *Coelastrum microporum*; *Merismopedia glauca*; *Pediastrum boryanum*; diatoms—*Navicula*, *Coscinodiscus*, etc.

Measurements.—The following is a series of measurements of Lost Lake examples:

No.	Date, 1908	MEASUREMENTS IN MM			Remarks
		Length	Alt.	Diam.	
1189	August 20	85	54	32	Fanshaped female.
1260	September 7	97.4	55	31	Fanshaped female, gravid.
1215	August 20	87	46	35.6	Fanshaped female.
1224	August 20	98	56	26	Fanshaped female.
1245	August 20	90	51	32.8	Fanshaped female.
1235	August 20	98	48.9	36.3	Male.
1188	August 20	102	53	36	Male.
1221	August 20	100	51	37	Male.
1223	August 20	96	51.4	34.8	Male.
1228	August 20	102.3	53.7	33	Male.

Most of these shells blistered posteriorly.  
 The males are fairly like those of river; the females are more fanshaped. Weight of the 10 shells, 15 oz.; only a few are rayed.

12. POCKETBOOK

LAMPSILIS VENTRICOSA (Barnes)

Rather common at the Long Point mussel bed; a few found in the bed by Farrar's and a few in Lost Lake. The species as found in the lake is markedly dwarfed and quite different in appearance from the usual river form. There are two types in the Long Point bed. One consisting of females, having the post basal inflation of the shell characteristic of that sex, not exactly as in the river form, however, but somewhat more restricted; this feature, along with a peculiar stain of the epidermis which conceals the normal coloring of the shell, causes them to very closely resemble a short female *L. luteola*. The other type, an oval shell without the post-basal inflation, was at first taken to represent the males, but some of them were found to contain glochidia. These, too, bear a marked resemblance to *L. luteola*, and the only way to distinguish the two species, as they occur in the lake, is by an examination of the umbonal sculpture. This in *ventricosa* consists of a few coarse parallel ridges; in *luteola* the sculpture is of numerous fine wavy lines.

The lake *L. ventricosa* was so markedly different from the species as usually known that it was compared with a large series of both lake and river forms. Of river shells only a few from the central part of the Maumee, where for some reason the shells are markedly dwarfed, bore any close resemblance to it. None was found in any of the neighboring lakes with which to compare them. In some of the small lakes of Michigan where Dr. Robert E. Coker collected, he experienced a similar difficulty in distinguishing *L. ventricosa* and *L. luteola*. He sent sets of criti-

cal specimens to Mr. Bryant Walker of Detroit, Mich., who identified the shells with a few coarse straight undulations on the beaks as *Lampsilis ventricosa canadensis* and the others as *L. luteola*.

The Maxinkuckee specimens were also compared with *L. ventricosa* from Lake Champlain, and were found to be much like them. The Champlain examples which were free from staining of the epidermis more closely resembled in color the *ventricosa* of the rivers.

The specimens of *L. ventricosa* differed considerably in the different beds. Lost Lake examples are usually rather small, and are stained a peculiar ashy-gray. Those from the near Farrar's are mostly small and apparently young and are rather well rayed; they resemble river forms more closely than any others in the lake.

The large oval *L. ventricosa* of Long Point are the heaviest shells of the lake. A peculiarity of several of these shells is a conspicuous rib-like thickening on the inside, extending from near the umbonal cavity postero-ventrally. The nacre is soft satiny in luster, and not very iridescent. This oval form of *ventricosa* found at Long Point furnishes the only shell in the lake that could be used to any advantage in the manufacture of buttons, and even it produces rather inferior material. Some of these shells were sent to a button factory at Davenport and buttons were made of them. The following is a set of measurements of these large shells:

No.	Date, 1907	Length mm.	Alt. mm.	Diam. mm.	Remarks
1	September 24	114	74.8	53	Female gravid.
2	October 30	107.6	65.5	54.8	
3	October 2	105.2	63.7	52.5	
4	October 30	92.5	60.4	53.7	Female gravid.
5	October 30	103.7	67.3	49.3	Dorsal baroques.
6	October 17	98.6	60.2	55.5	Arcuate; baroque found.
7	October 20	101.7	63.6	52.2	
8	October 30	94.6	55.4	52.2	Nacre diseased and blistered.
9	October 17	95.6	55.7	49	
10	October 17	91.5	60.4	49.5	

Although the reproductive phase of *L. ventricosa* of the lake is much less conspicuous than in the river mussels, most of them apparently succeed in reproducing themselves. Most of the females found later in autumn have more or less numerous glochidia in the gills. No infected fishes or very young mussels of this species were seen.

The most common parasite is *Atax*, and it is not particularly abundant. Of six examples collected near Farrar's July 24, 1909,

the first contained 9 of the mites, the second 4, the third 15, with *Atax* eggs in the mantle and body, the fourth 12 *Atax* and numerous eggs of the mite on the inner surface of the mantle, the fifth 3 *Atax* with eggs and the sixth 7 *Atax* with eggs and egg scars. No other parasites were noted. No pearls were found, only a few irregular slugs.

In 1906 some of the immense *L. ventricosa* of Yellow River were planted in the lake near shore not far from the old ice office. A few died shortly after planting but near the same place 2 years later some of the mussels were found alive and apparently thriving. Two of the large females were killed and examined. Altho this was at a time when this species is usually gravid, one of these individuals was sterile, apparently having failed to become impregnated. The influence of its residence in the lake was marked by a dark stain which covered the exposed portion of the shell. The other had a few eggs in the gills, and numerous marginal cysts in the mantle. About 10 *Atax* among the gills, and numerous distomids on the outside surface of the mantle in the umbonal cavity.

### 13. LAMPASILIS MULTIRADIATA (Lea)

Not abundant in the lake; occasional shells are found along shore, and now and then they are encountered in the piles of shells where muskrats have been feeding. A few living examples were found in the mussel bed near the mouth of Norris Inlet and a few at Long Point bed. In all hardly a dozen living examples were secured; of 563 shells taken from a pile left by a muskrat at Long Point in 1907, only 1 was of this species. This mussel, as it occurs in the lake, is not nearly so attractive as river specimens, being dwarfed and so deeply stained that the rays are inconspicuous, being usually black or dull brown instead of green.

This species was found in unusual abundance in the Tippecanoe River at Delong, and a considerable number was observed spawning during the autumn of 1908. While spawning, this mussel is a very conspicuous spectacle. It lies either on its back, or more usually with the posterior end directly upward, and the showy edges of the mantle, which are of a yellowish brown color, and cross-barred with narrow lines which are continuous with the fine rays of the epidermis, look a good deal like a small darter lying on the bottom. Long waving pennant-like flaps, with a showy black spot at the base of each are developed, and this portion of the mussel is made still more conspicuous by reason of periodic violent spasmodic contractions.

At the Tippecanoe River this is one of the favorite foods of the muskrat, and it must be difficult for them to hold their own against that rodent.

14. MICROMYA FABALIS (Lea)

Rare; previous to 1913 only one shell had been found; this was picked up on the north shore of the lake in 1907. In 1913 several shells, recently cleaned out by some animal, probably a muskrat, were found at the wagon bridge. This species is fairly common in Tippecanoe Lake and frequent in the Tippecanoe River at DeLong where it was collected in shallow water near shore in rather stiff blue clay. It is the smallest of our Unionidæ. The white or bluish white nacre has an exceedingly brilliant luster.

Several other species of mussels have been recorded for the lake, among them *Quadrula lachrymosa* (Lea), *Alasmodonta marginata* Say, *Symphynota compressa* Lea, *Anodontoides ferussacianus* (Lea), *Ptychobranhus phaseolus* (Hildreth), *Obovaria circulus* Lea, *Lampsilis parva* (Barnes), and *Lampsilis gracilis* (Barnes). We have seen representatives of none of these species from the lake, and while some, such as *A. ferussacianus*, are very probably present, others are very improbable.

MOLLUSKS OTHER THAN UNIONIDÆ

About 116 species of mollusks in addition to the Unionidæ are known to occur in Lake Maxinkuckee or its immediate vicinity. Specimens of nearly all of these species were collected during our investigations; others were collected by the late L. E. Daniels. These collections were referred to Dr. Paul Bartsch of the United States National Museum for study and report. Other duties have prevented Dr. Bartsch from completing his full report on these mollusks. He has, however, supplied the following list of his identifications. This shows that the Lake Maxinkuckee molluscous fauna is a remarkably rich one, the total number of species, including the Unionidæ, being not fewer than 130.

FRESHWATER MOLLUSKS

1. *Lymnæa palustris* (Müller)
2. *Lymnæa obrussa exigua* (Lea)
3. *Lymnæa danielsi* Baker
4. *Lymnæa dalli* Baker
5. *Lymnæa humilis* (Say)
6. *Lymnæa desidiosa* (Say)
7. *Planorbis bicarinatus* Say