

## Host Identification for *Strophitus undulatus* (Bivalvia: Unionidae), the Creeper, in the Upper Susquehanna River Basin, Pennsylvania

ELLEN VAN SNIK GRAY,<sup>1</sup> WILLIAM A. LELLIS, JEFFREY C. COLE  
AND CONNIE S. JOHNSON

U.S. Geological Survey, Northern Appalachian Research Laboratory, Wellsboro, Pennsylvania 16901

**ABSTRACT.**—Hosts for *Strophitus undulatus* (Bivalvia: Unionidae) were identified through laboratory infestations. *Strophitus undulatus* had a low degree of host specificity, transforming on 15 of 22 species examined, including three non-native species and one anadromous species. Suitable hosts included five cyprinid species, two salmonids, two centrarchids, two percids and *Acipenser oxyrinchus* (Atlantic sturgeon), *Ameiurus natalis* (yellow bullhead), *Cottus cognatus* (slimy sculpin) and *Notophthalmus viridescens viridescens* (red-spotted newt). Ten hosts were previously unknown. No metamorphosis was observed in a control treatment without a potential host. Recovery of juveniles occurred 12–41 d after infestation at 13–18 C. Duration of glochidial attachment to the hosts declined with increasing water temperature.

### INTRODUCTION

Freshwater mussels of the families Margaritiferidae and Unionidae are the most imple North American aquatic group, with only 24% of 297 species considered stable (Bog 1993; Williams *et al.*, 1993). Conservation efforts for this group continue to be hampered by lack of specific life history information, particularly knowledge of host relationships. Freshwater mussels have a highly specialized life history in which larvae (glochidia), bred by females in modified gill chambers (marsupia) and released when mature, undergo a brief period as obligate ectoparasites on fish or amphibians. This life cycle is followed by rare exceptions (Kondo, 1990; Lellis and King, 1998). Host specificity varies widely across species however, with some unionids parasitizing a wide variety of unrelated host species (Trdan and Hoeh, 1982; Hove *et al.*, 1997), and others using only a few host species, of closely related (Yeager and Neves, 1986; Michaelson and Neves, 1995). Inappropriate hosts shed attached glochidia, usually within a few days, whereas appropriate hosts carry glochidia until transformation into the juvenile form, which excysts and becomes free-living. However, for only one quarter of the North American mussel species have been reported (Hogga 1992; Watters, 1994). We identified hosts of *Strophitus undulatus*, the creeper, in the upper Susquehanna River basin, Pennsylvania, through laboratory studies. Despite early reports that *S. undulatus* transforms without the use of a host (Lefevre and Curtis, 1911; How 1914a), subsequent studies have shown *S. undulatus* uses fish hosts (Baker, 1928; How 1914a, 1997; Watters *et al.*, 1998; Wicklow and Beisheim, 1998).

### METHODS

Laboratory experiments were conducted to identify suitable hosts for *Strophitus undulatus*. When possible, potential hosts were collected from areas devoid of mussels to av

<sup>1</sup> Corresponding author, Present address: U.S. National Park Service, Center for Urban Ecology, 4 MacArthur Boulevard, N.W., Washington, DC 20007; Telephone (202) 342-1443 extension 223; (202) 282-1031; e-mail: ellen\_gray@nps.gov

using fish with acquired immunity to glochidia (Arey, 1923). Most fishes were collected from Pine Creek, Tioga County, Pennsylvania, a tributary to the West Branch of the Susquehanna River. *Cottus cognatus* (slimy sculpin) were collected from Straight Run, and *Etheostoma zonale* (banded darter) from Little Pine Creek, both tributaries to Pine Creek. *Bufo americanus* tadpoles (American toad) and adult *Notophthalmus viridescens viridescens* (red-spotted newt) were collected from local ponds. *Acipenser oxyrinchus* (Atlantic sturgeon), *Salmo salar* (Atlantic salmon) and *Salvelinus fontinalis* (brook charr), were taken from captive bred populations at the USGS Northern Appalachian Research Laboratory in Wellsboro, Pennsylvania. *Ameiurus natalis* (yellow bullhead), *Micropterus salmoides* (large mouth bass) and *Perca flavescens* (yellow perch) were collected from Tamarack Lake, Crawford County, Pennsylvania. We attempted to expose *S. undulatus* glochidia to a variety of aquatic species, including those likely to be present during glochidial release by *S. undulatus* as well as several non-native species (*Oncorhynchus mykiss* (rainbow trout), *Ambloplites rupestris* (rock bass), *M. salmoides*, *E. zonale*) and anadromous species (*A. oxyrinchus*, *S. salar*) that may have been present historically (Cooper, 1983; Lee *et al.*, 1980).

A total of 86 *Strophitus undulatus* ranging in size from 41–80 mm and 8–50 g (mean 65 ± 8 mm, 27 ± 9 g) were collected from Pine Creek, Tioga County, Pennsylvania (41°44.099'N, 077°25.857'W) between 15 April and 13 May 1999 and transported in cooler without water to the USGS Northern Appalachian Research Laboratory in Wellsboro, Pennsylvania for experimentation. Unionid nomenclature follows Turgeon *et al.* (1988) and voucher specimens were deposited in the American Museum of Natural History, New York. At the laboratory, mussels were cleaned and measured, then divided equally into two glass aquaria (122 × 46 × 51 cm, L × W × D) containing 10 cm of sand substrate. Aquaria were supplied with 1 liter/min well water (9.8 C ambient) heated or chilled with a submerged stainless steel heat-exchanger connected to an external refrigerated circulator (Model 675 Fisher Scientific, Pittsburgh, Pennsylvania). Water was circulated within each aquarium by aeration from a 31 cm airstone set opposite the drain end of the tank. Each aquarium was illuminated with two 40W fluorescent lamps (Cool White Deluxe, Osram Sylvania, Inc. Versailles, Kentucky) set 50 cm above the water surface and controlled by a programmable digital timer (Model 810030, Sper Scientific, Ltd., Scottsdale, Arizona). Photoperiod and temperature were systematically adjusted in one aquarium to match 26 y average condition for Pine Creek at Cedar Run, Pennsylvania (1944–1970 period of record; Flippo, 1975). In the second aquarium, photoperiod and temperature were held constant at average 20 April conditions (8 C, 13.5 h light) for 4 wk before resuming normal seasonal patterns to delay glochidial release. Mussels were fed twice daily a mixture of cultured microalgae that included *Nanochloropsis* sp. (Florida Aqua Farms, Inc., Dade City, Florida) and *Neochloris oleoabundans*, *Bracteacoccus grandis* and *Bracteacoccus giganteus* (University of Texas, Austin Texas).

Naturally released glochidia and conglomerates were removed with a pipette from the *Strophitus undulatus* aquaria for use in laboratory infections. Glochidia were considered viable if they snapped shut upon exposure to salt (Zale and Neves, 1982). Potential hosts were examined for existing glochidial infections prior to experimentation. Small fishes tadpoles and *Notophthalmus viridescens* (1–24 individuals/species) were exposed to *S. undulatus* glochidia in a 600 ml beaker aerated with an airstone at 12–19 C. Larger fishes (2–5 individuals/species; *Acipenser oxyrinchus*, *Ameiurus natalis*, *Salmo salar*, *Salvelinus fontinalis*, *Micropterus salmoides*, *Perca flavescens*) were exposed to glochidia in buckets aerated by an airstone at 13–15 C. Glochidia were maintained in suspension by frequent agitation. Duration of exposure varied from 15–75 min, depending on the size of the container and the susceptibility of the host to glochidial attachment. Potential hosts were placed in aer

ated, flow-through, 38–76 liters aquaria by species at 13–18 C and fed chopped squid, fi or frozen brine shrimp daily. Uneaten food was removed promptly. No substrate was plac in the tanks, however potential hosts were provided with PVC pipe halves (10 cm long 5 cm wide  $\times$  2.5 cm high) for cover and newts were provided with a small board held ju below the water surface. We conducted only one replicate for each species due to spat limitations. A control treatment was included in which glochidia were added to a 76-lit aquarium in the absence of a potential host in order to determine if transformation cou occur without the use of a host. In addition, 50–100 conglutinates were placed in each 27 sterile plastic petri dishes (100  $\times$  15 mm, Plasta-Medic, Carson, California) containi either distilled water ( $n = 13$ ) or boiled well water ( $n = 14$ ). Dishes were cultured in incubator (Model 5430, Precision Scientific, Chicago, Illinois) at  $10 \pm 1$  C for 8–49 d an checked periodically for viability and development.

Glochidial attachment was verified for each species 2 d after exposure to glochidia. Af a species carried glochidia for 4 d, we inspected material on the bottom of the tank thr times a week for glochidia and transformed juveniles. The tank bottom was siphoned in a 100  $\mu\text{m}$  mesh sieve, washed into a gridded petri dish and examined under a microscop Juveniles were distinguished from glochidia by the presence of a foot, and were usua crawling on the bottom of the petri dish or in debris. Number of glochidia and juveni were tabulated and a subsample preserved. Siphoning continued for at least 10 d after t last juveniles were found. A species was considered a host if encystment and metamorpho to the juvenile stage occurred.

## RESULTS

*Strophitus undulatus* released glochidia 26 April to 2 June (8.5–16.9 C) when held at photoperiod and temperature regime that mimicked average natural conditions for Pi Creek (Fig. 1). The majority of conglutinates were released between 12 May and 27 M (13.6–15.6 C). Mussels subjected to a 4 wk seasonal delay released glochidia from 4 May 18 June (8.0–15.6 C) with the majority released from 26 May to 18 June (13.3–15.6 C). There were two large release events within the delayed season tank on 9 June (15.5 C) an 18 June (15.6 C). We could not determine how many mussels contributed to any sing release. Laboratory infestations were begun on 26 April 1999 and were completed by July 1999.

*Strophitus undulatus* successfully metamorphosed on 15 of 22 potential hosts, includi five cyprinid species, two salmonids, two centrarchids and two percids, as well as *Acipen. oxyrhynchus*, *Ameiurus natalis*, *Cottus cognatus* and *Notophthalmus viridescens* (Table 1). Three host species are non-native to the Susquehanna River drainage: *Oncorhynchus myki*, *Ambloplites rupestris* and *Micropterus salmoides*, while one anadromous species, *A. oxyrhy chus*, also acted as a host (Table 1). Hosts produced 1–36 juveniles with transformation rat varying from 2–51% (Table 1). The weighted average number of days until metamorpho and mean temperature were related. Duration of glochidial attachment to the hosts c lined with increasing temperature (Fig. 2). Seven species did not act as hosts, includi two cyprinid species, one sucker, one ictalurid, one salmonid and one percid (Table 1). transformation of glochidia was observed in the control treatment. Glochidia incubated petri dishes remained viable for as long as 9 d postrelease, but no development was observ among individuals that were either free, attached to, or contained within conglutinat. Conglutinate stability varied, but some held in distilled water remained intact in excess 14 d.




FIG. 1.—Release of glochidia by *Strophitus undulatus* held in tanks under a natural photoperiod at temperature regime ('natural conditions') or a 'delayed season,' in which temperature and photoperiod were held constant for 4 wk at 8 C, 13.5 h light before resuming normal seasonal patterns. Release events are indicated by diamonds for natural conditions and by triangles for delayed seasonal conditions.

#### DISCUSSION

We identified 15 hosts for *Strophitus undulatus*, ten of which were previously unknown bringing the total number of reported hosts to 29 for this species (Baker, 1928; Hove *et al.* 1997; Watters *et al.*, 1998; Wicklow and Beisheim, 1998; Wicklow, pers. comm.—yellow perch). *Acipenser oxyrinchus*, *Campostoma anomalum*, *Luxilus cornutus*, *Nocomis biguttatus*, *Rhinichthys atratulus*, *Oncorhynchus mykiss*, *Salvelinus fontinalis*, *Ambloplites rupestris*, *Etheostoma olmstedii* and *Notophthalmus viridescens* are all reported as hosts for *S. undulatus* for the first time. Two species that have been previously identified as hosts for *S. undulatus*—*Etheostoma zonale* and *Pimephales notatus*, did not metamorphose glochidia in this study. *Pimephales notatus* shed all attached glochidia in the first 8 d. *Etheostoma zonale* shed no glochidia in the first 10 d, however after a period of inactivity, shed two more glochidia after 20 d and one more on day 29. As reported by Watters and O'Dee (1996), this may represent acquired immunity, although further studies are needed to test this hypothesis. Early reports that *S. undulatus* did not require a host (Lefevre and Curtis, 1911; Howar 1914a) were not corroborated by this study. Duration of metamorphosis and temperature were significantly related in this study and others, with fewer days until metamorphosis occurring at higher temperatures (Howard and Anson, 1922; Zale and Neves, 1982; Weaver *et al.*, 1991). In addition, there was a trend towards a shorter interval of metamorphosis at higher temperatures.

Hoggarth (1992) summarized known host relationships for unionids and his data revealed

TABLE 1.—Results of laboratory trials to determine hosts of *Strophitus undulatus*. Non-native species are designated with an asterisk

Potential host species	Number of individuals exposed	Mean length (mm)	Mean temp. (C)	Number of glochidia shed	Period of metamorphosis (days)	Number of juveniles recovered	Percent metamorphosed
Control	n/a	n/a	18.1	102	n/a	0	0%
<i>Aeroperca oxyrinchus</i> , Atlantic Sturgeon	2	210	14.9	3	17	1	25%
<i>Camptostoma anomatum</i> , Central stoneroller	1	81	15.5	7	20	2	22%
<i>Exoglossum maxillingua</i> , Cutlips minnow	7	65	16.9	8	n/a	0	0%
<i>Luxilus cornutus</i> , Common shiner	7	93	16.8	35	13-22	36	51%
<i>Nocomis biguttatus</i> , River chub	3	77	16.9	57	12-16	5	8%
<i>Pimephales notatus</i> , Bluntnose minnow	6	51	15.7	7	n/a	0	0%
<i>Rhinichthys atratulus</i> , Blacknose dace	1	54	13.4	3	29	1	25%
<i>Rhinichthys cataractae</i> , Longnose dace	2	82	15.6	7	17-22	7	50%
<i>Catostomus commersoni</i> , White sucker	8	97	15.5	7	n/a	0	0%
<i>Ameiurus natalis</i> , Yellow bullhead	4	177	16.3	14	17-18	2	13%
<i>Naturus insignis</i> , Margined madtom	7	108	13.5	100	n/a	0	0%
<i>Oncorhynchus mykiss</i> *, Rainbow trout	1	92	13.8	84	14-19	2	2%
<i>Salmo salar</i> , Atlantic salmon	4	97	13.4	5	n/a	0	0%
<i>Sabotinus fontinalis</i> , Brook charr	5	85	15.5	11	16-21	3	21%
<i>Cottus cognatus</i> , Slimy sculpin	7	54	13.4	33	34	1	3%
<i>Ambloplites rupestris</i> *, Rock bass	4	70	13.8	69	18-29	9	12%
<i>Micropterus salmoides</i> *, Largemouth bass	3	151	13.3	76	26	4	5%
<i>Etheostoma olmstedii</i> , Tesselated darter	6	60	13.2	94	32-37	20	18%
<i>Etheostoma zonale</i> *, Banded darter	4	50	13.2	16	n/a	0	0%
<i>Perca flavescens</i> , Yellow perch	3	133	13.3	149	20-41	27	15%
<i>Bufo americanus</i> , American toad	24	20	18.4	14	n/a	0	0%
<i>Notophthalmus viridescens</i> , red-spotted newt	6	68	14.9	4	21	1	20%




FIG. 2.—Relationship between weighted average number of days until metamorphosis and temperature for *Strophitus undulatus*. Each point represents an identified host for *S. undulatus*.

a mean number of 4.4 host/species. *Strophitus undulatus* is therefore a host generalist, using more than the average number of taxonomically diverse host species. Only *Pyganodon grandis*, the giant floater, has more reported hosts (32) (Hoggarth, 1992). In addition to *S. undulatus* and *P. grandis*, many other species in the subfamily Anodontinae are also host generalists (Trdan and Hoeh, 1982; Barnhart and Roberts, 1997; Haag and Warren, 1997; Watters and O'Dee, 1998). Use of such a large number of host species should protect these unionid species against population decline as well as facilitate dispersal and recolonization (Trdan and Hoeh, 1982; Kat, 1984).

Host species differed significantly in attachment and metamorphosis rates, with 4 to 17% of glochidia attaching to the hosts, and 2 to 51% of attached glochidia becoming juveniles. Many factors may affect variation in attachment, including immune response, swimming behavior and orientation of the host, although the role of these factors in susceptibility to attachment is unknown. Studies of other mussel species have also found a range of suitability among identified hosts (Waller and Holland-Bartels, 1988; Haag and Warren, 1997; Watter and O'Dee, 1998). Attachment rates in the field are typically significantly less than those obtained in the laboratory (Zale and Neves, 1982; Weaver *et al.*, 1991; Bruenderman and Neves, 1993); therefore percent metamorphosed is an important variable in the reproductive success of unionid species. We hypothesized that species with metamorphosis rate below 10% may be marginally suitable hosts for *Strophitus undulatus*, including *Nocomis micropogon*, *Oncorhynchus mykiss*, *Cottus cognatus* and *Micropterus salmoides*. Future studies should examine the importance of these hosts in the field. Nevertheless, loss of these host is presumed to have vastly different effects upon *S. undulatus* populations as compared to

species such as *Luxilus cornutus*, *Rhinichthys cataractae*, *Etheostoma olmstedii* or *Perca flavescens*, which had metamorphosis rates of 15–51%.

*Strophitus undulatus* was also able to use both non-native and anadromous fishes as hosts including *Acipenser oxyrinchus*, *Oncorhynchus mykiss*, *Ambloplites rupestris* and *Micropterus salmoides*. *Oncorhynchus mykiss*, *A. rupestris* and *M. salmoides* are present in Pine Creek and could act as hosts in the upper Susquehanna basin. In areas in which *M. salmoides* is not present it is also used as a host. Sturgeon are used as hosts for three other unionids: *Lampsilis teres*, the yellow sandshell, *Obovaria olivaria*, the hickorynut and *Quadrula pustulosa*, the blue mussel, all of which use the shovelnose sturgeon, *Scaphirhynchus platyrhynchus*, although transformation was not observed for *L. teres* or *Q. pustulosa* (Surber, 1913; Howard, 1951; Wilson, 1916; Coker *et al.*, 1921). Dams and other barriers to upstream movements drastically affect the distribution of unionids with anadromous fish as primary hosts, for example in *Anodonta imbecilla*, the alewife floater, and its host, *Alosa pseudoharengus*, the alewife (Davenport and Warmuth, 1965; Smith, 1985).

Identification of mussel hosts has traditionally focused on fish species, however recent studies have shown that amphibians act as hosts for unionids as well (Howard, 1951; Hines, 1941; Walker, 1981; Watters, 1997; Watters and O'Dee, 1998; Wicklow and Beishwiler, 1998). As recommended by Watters (1997), future studies need to expose a wider variety of aquatic organisms to mussel glochidia, rather than focus solely on fishes. This is the case of a unionid capable of using newts as hosts. The red-spotted newt is unusual in that its juveniles are completely terrestrial, whereas both the larval and adult stages are aquatic (Conant and Collins, 1991). Watters and O'Dee (1998) attempted infestations of *Lampsilis cardium* and *Utterbackia imbecilla* with decapod crustaceans but no juveniles were produced.

*Strophitus undulatus* is a habitat generalist, and uses two strategies for targeting potential hosts: first conglutinates are released into the water column that are attacked by fishes, a potential prey item, second, the conglutinates degrade, leaving glochidia on the stream bottom that can attach to fins and other soft tissues of benthic species (Wicklow and Beishwiler, 1998). Thus, characteristics of *S. undulatus* glochidial release and habitat use complement its ability to parasitize a wide variety of aquatic species. Although host studies conducted in the laboratory do not unequivocally determine hosts used in the field, correlation of ecological characteristics of potential hosts such as feeding mode, habitat, spawning timing and habitats used in spawning can indicate which potential hosts are most likely to encounter mussel glochidia in situ. In Pine Creek *S. undulatus* generally releases glochidia in April to June in runs or pools. Using this information, we predict that species that are found on small drifting items in the water column or items on the substrate, use these habitats during spring months, and transform 10% or more *S. undulatus* glochidia would most likely be hosts of *S. undulatus* in Pine Creek. Using these criteria, we predict that *Camptostoma omalum*, *Luxilus cornutus*, *Rhinichthys atratulus*, *Rhinichthys cataractae*, and *Etheostoma olmstedii* are most likely hosts used by *S. undulatus* in Pine Creek and should be targeted in future field studies.

*Acknowledgments.*—We thank G. Ayers, D. S. Dropkin, R. L. Flynn, M. J. Gray and L. A. Ridell for assistance in the field and laboratory. We thank M. Marcinko, L. Hines and the Tionesta Fish Hatchery of the Pennsylvania Fish and Boat Commission for providing fish for the study. Rita Vilella and I. Smith reviewed the manuscript.

#### LITERATURE CITED

- AREY, L. B. 1923. Observations on an acquired immunity to a metazoan parasite. *J. Exp. Zool.*, 38: 381.

- BAKER, F. C. 1928. The fresh water Mollusca of Wisconsin. Part II. Pelecypoda. *Bull. Wisc. Geol. Nat. Hist. Surv.* No. 70:1-495.
- BARNHART, M. C. AND A. ROBERTS. 1997. Reproduction and fish hosts of unionids from the Ozark uplifts, p. 16-20. *In*: K. S. Cummings, A. C. Buchanan, C. A. Mayer and T. J. Naimo (eds.) Conservation and management of freshwater mussels II: initiatives for the future: Proceeding of an Upper Mississippi River Conservation Committee symposium, St. Louis, Missouri, October 1995. Upper Mississippi River Conservation Committee, Rock Island, Illinois.
- BOGAN, A. E. 1993. Freshwater bivalve extinctions (Mollusca: Unionoida): a search for causes. *Am. Zool.*, 33:599-609.
- BRUENDERMAN, S. A. AND R. J. NEVES. 1993. Life history of the endangered fine-rayed pigtoe *Fuscomai cuneobus* (Bivalvia: Unionidae) in the Clinch River, Virginia. *Am. Malacological Bull.*, 10:83-91
- COKER, R. E., A. F. SHIRA, H. W. CLARK AND A. D. HOWARD. 1921. Natural history and propagation of fresh-water mussels. *Bulletin of the U.S. Bureau of Fisheries*, 37:77-181.
- CONANT, R. AND J. T. COLLINS. 1991. A field guide to reptiles and amphibians. Houghton Mifflin Company, Boston. 450 p.
- COOPER, E. L. 1983. Fishes of Pennsylvania and the Northeastern United States. The Pennsylvania State University Press, University Park, Pennsylvania. 243 p.
- DAVENPORT, D. AND M. WARMUTH. 1965. Notes on the relationship between the freshwater mussel *Anodonta implicata* Say and the alewife *Pomolobus pseudoharengus* (Wilson). *Limnol. and Oceanog.*, 10:R74-R78.
- FLIPPO, H. N., JR. 1975. Temperatures of streams and selected reservoirs in Pennsylvania. *Bulletin No. 11, Department of Environmental Resources, Commonwealth of Pennsylvania, Harrisburg*. 95 p.
- HAAG, W. R. AND M. L. WARREN, JR. 1997. Host fishes and reproductive biology of 6 freshwater mussel species from the Mobile basin, USA. *J. N. Am. Benthol. Soc.*, 16:576-585.
- HOGGARTH, M. A. 1992. An examination of the glochidia—host relationships reported in the literature for North American species of Unionacea (Mollusca: Bivalvia). *Malacology Data Net*, 3:1-30.
- HOVE, M. C., R. A. ENGELKING, M. E. PETELER, E. M. PETERSON, A. R. KAPUSCINSKI, L. A. SOVELL AND E. R. EVERS. 1997. Suitable fish hosts for glochidia of four freshwater mussels, p. 21-25. *In*: K. S. Cummings, A. C. Buchanan, C. A. Mayer and T. J. Naimo (eds.). Conservation and management of freshwater mussels II: initiatives for the future: Proceedings of an Upper Mississippi River Conservation Committee symposium, St. Louis, Missouri, October 1995. Upper Mississippi River Conservation Committee, Rock Island, Illinois.
- HOWARD, A. D. 1914a. A second case of metamorphosis without parasitism in the Unionidae. *Science* 40:353-355.
- . 1914b. Some cases of narrowly restricted parasitism among commercial species of fresh water mussels. *Trans. Am. Fish. Soc.*, 44:41-44.
- . 1951. A river mussel parasitic on a salamander. *Nat. Hist. Misc.*, 77:1-6.
- , AND B. J. ANSON. 1922. Phases in the parasitism of the Unionidae. *J. Parasitol.*, 9:70-84.
- KAT, P. W. 1984. Parasitism and the Unionacea (Bivalvia). *Biolog. Rev.*, 59:189-207.
- KONDO, T. 1990. Reproductive biology of a small bivalve *Grandidiera burtoni* in Lake Tanganyika. *Venus* 49:120-125.
- LEE, D. S., C. R. GILBERT, C. H. HOCUTT, R. E. JENKINS, D. E. MCALLISTER AND J. R. STAUFFER, JR. 1986. Atlas of North American freshwater fishes. North Carolina State Museum of Natural History 867 p.
- LEFEVRE, G. AND W. C. CURTIS. 1911. Metamorphosis without parasitism in the Unionidae. *Science*, 33 863-865.
- LELLIS, W. A. AND T. M. KING. 1998. Release of transformed juveniles by the green floater, *Lasmigon subviridis*. *Triannual Unionid Report*, 16:23.
- MICHAELSON, D. L. AND R. J. NEVES. 1995. Life history and habitat of the endangered dwarf wedge mussel *Alasmidonta heterodon* (Bivalvia: Unionidae). *J. N. Am. Benthol. Soc.*, 14:324-340.
- SMITH, D. G. 1985. Recent range expansion of the freshwater mussel *Anodonta implicata* and its relationship to clupeid fish restoration in the Connecticut River system. *Freshwat. Invertebr. Biol.* 4:105-108.



- SESHAIYA, R. V. 1941. Tadpoles as hosts for the glochidia of the fresh-water mussel. *Current Science*, 535-536.
- SURBER, T. 1913. Notes on the natural hosts of fresh-water mussels. *Bulletin of the U.S. Bureau of Fishes* 32:110-116.
- TURGEON, D. D., A. E. BOGAN, C. V. COAN, W. K. EMERSON, W. G. LYONS, W. L. PRATT, C. F. E. ROE, G. ROSENBERG, A. SCHELTEMA, F. G. THOMPSON AND J. D. WILLIAMS. 1988. Common and entific names of aquatic invertebrates from the United States and Canada: Mollusks. *Am. J. Soc. Special Publication* 16. 227 p.
- TRDAN, R. J. AND W. R. HOEH. 1982. Eurytopic host use by two congeneric species of freshwater mu (Pelecypoda: Unionidae: Anodonta). *Am. Midl. Nat.*, 108:381-388.
- WALKER, K. F. 1981. The ecology of freshwater mussels in the River Murray. *Austr. Water. Res. 7 Pap.*, 63:1-110.
- WALLER, D. L. AND L. E. HOLLAND-BARTELS. 1988. Fish hosts for glochidia of the endangered freshw mussel *Lampsilis higginsii* Lea (Bivalvia: Unionidae). *Malacolog. Rev.*, 21:119-122.
- WATTERS, G. T. 1994. An annotated bibliography of the reproduction and propagation of the Un oidea (Primarily of North America). *Ohio Biological Survey Miscellaneous Contributions* No 158 p.
- . 1997. Glochidial metamorphosis of the freshwater mussel *Lampsilis cardium* (Bivalvia: Uni dae) on larval tiger salamanders, *Ambystoma tigrinum* ssp. (Amphibia: Ambystomidae). *Ca Zool.*, 75:505-508.
- , AND S. H. O'DEE. 1996. Shedding of untransformed glochidia by fishes parasitized by *Lamp fasciola* Rafinesque, 1820 (Mollusca: Bivalvia: Unionidae): evidence of acquired immunity in field? *J. Freshw. Ecol.*, 11:383-389.
- , AND S. H. O'DEE. 1998. Metamorphosis of freshwater mussel glochidia (Bivalvia: Unionic on amphibians and exotic fishes. *Am. Midl. Nat.*, 139:49-57.
- , S. H. O'DEE AND S. CHORDAS. 1998. New potential hosts. *Triannual Unionid Report*, 15:27-28.
- WEAVER, L. R., G. B. PARDUE AND R. J. NEVES. 1991. Reproductive biology and fish hosts of the T nnessee Clubshell *Pleurobema oviforme* (Mollusca: Unionidae) in Virginia. *Am. Midl. Nat.*, 126:82-89.
- WICKLOW, B. J. AND P. M. BEISHEIM. 1998. Life history studies of the squawfoot mussel *Strophitus dulatus* in the Piscataquog River watershed, New Hampshire, p. 41. *In: Conservation, Cap Care, and Propagation Freshwater Mussel Symposium*, Columbus, Ohio, March 1998.
- WILLIAMS, J. D., M. L. WARREN, JR., K. S. CUMMINGS, J. L. HARRIS AND R. J. NEVES. 1993. Conserva status of freshwater mussels of the United States and Canada. *Fisheries*, 18:6-22.
- WILSON, C. B. 1916. Copepod parasites of fresh-water fishes and their economic relations to mu glochidia. *Bulletin of the U.S. Bureau of Fisheries Document Number* 824. 374 p.
- YEAGER, B. L. AND R. J. NEVES. 1986. Reproductive cycle and fish hosts of the Rabbit's Foot mu: *Quadrula cylindrica strigillata* (Mollusca: Unionidae) in the Upper Tennessee River drain. *Am. Midl. Nat.*, 116:329-340.
- ZALE, A. V. AND R. J. NEVES. 1982. Fish hosts of four species of lampsiline mussels (Mollusca: Unionic in Big Moccasin Creek, Virginia. *Can. J. Zool.*, 60:2535-2542.

